



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2018

Does orthodontic treatment have a permanent effect on tooth color? : A systematic review and meta-analysis

Kamber, Rita ; Papageorgiou, Spyridon N ; Eliades, Theodore

Abstract: **OBJECTIVES** Aim of this systematic review was to assess the effect of orthodontic treatment with fixed appliances on the tooth color of patients. **METHODS** Nine databases were searched up to May 2017 for clinical cohort studies on the effect of fixed appliance treatment on tooth color. After elimination of duplicate studies, data extraction, and risk of bias assessment according to the Cochrane guidelines, random effects meta-analyses of mean differences (MD) or means and their 95% confidence intervals (CIs) were performed, followed by GRADE (Grading of Recommendations Assessment, Development and Evaluation) assessment of the quality of evidence. **RESULTS** Three nonrandomized and one randomized study with a total of 138 patients (46% male, 54% female) with average age of 15.7 years were included. Tooth color of treated patients was significantly altered during or after orthodontic treatment (4 studies; average of 3.2 ΔE units; 95% CI = 2.0-4.4 ΔE units), which was more than the variation among controls (1 study; MD = 1.9 ΔE units; 95% CI = 1.7-2.2 ΔE units). However, the quality of evidence was very low, due to the inclusion of nonrandomized studies, bias, and imprecision. Re-analysis of raw study data indicated that significant differences in clinically discernable treatment-induced color changes were seen between chemically and light-cured adhesives and among the various tooth categories. **CONCLUSION** Existing evidence of very low quality indicates that orthodontic treatment might be associated with alterations of tooth color, which are however not consistently clinically discernible. Treatment-induced color alterations might be dependent on bonding material and tooth type, but evidence supporting this is weak.

DOI: <https://doi.org/10.1007/s00056-018-0123-7>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-169508>

Journal Article

Accepted Version

Originally published at:

Kamber, Rita; Papageorgiou, Spyridon N; Eliades, Theodore (2018). Does orthodontic treatment have a permanent effect on tooth color? : A systematic review and meta-analysis. *Journal of orofacial orthopedics = Fortschritte der Kieferorthopädie*, 79(2):73-82.

DOI: <https://doi.org/10.1007/s00056-018-0123-7>

Title Page

Does orthodontic treatment have a permanent effect on tooth color? A systematic review and meta-analysis

Rita Kamber¹, Spyridon N. Papageorgiou¹, Theodore Eliades¹

¹Clinic of Orthodontics and Pediatric Dentistry, Center of Dental Medicine, University of Zurich, Plattenstrasse 11, Zurich 8032, Switzerland

ORCIDs

S.N. Papageorgiou: 0000-0003-1968-3326

T. Eliades: 0000-0003-2313-4979

Short title: Orthodontics treatment and tooth color

Corresponding author: Spyridon N. Papageorgiou; Clinic of Orthodontics and Pediatric Dentistry, Center of Dental Medicine, University of Zurich, Plattenstrasse 11, CH-8032, Zurich; E: snpapage@gmail.com.

Words in abstract: 246

Words in text: 2651

Registration: PROSPERO (CRD42017064862)

Key words: orthodontics, orthodontic appliances, tooth color, adverse effects, clinical trials, systematic review, meta-analysis

Acknowledgements: We thank Dr. A. Karamouzou (private practice, Chios, Greece) for graciously providing the full dataset of their included study.

Conflicts of interest: None.

Does orthodontic treatment have a permanent effect on tooth color? A systematic review and meta-analysis

Abstract

Objectives Aim of this systematic review was to assess the effect of orthodontic treatment with fixed appliances on the tooth color of human patients.

Methods Nine databases were searched up to May 2017 for clinical cohort studies on the effect of fixed appliance treatment on tooth color. After elimination of duplicate studies, data extraction, and risk of bias assessment according to the Cochrane guidelines, random effects meta-analyses of Mean Differences (MD) or means and their 95% confidence intervals (CIs) were performed, followed by GRADE assessment of the quality of evidence.

Results Three non-randomized and one randomized study with a total of 138 patients (46% male/ 54% female) with average age of 15.7 years were included. Tooth color of treated patients was significantly altered during or after orthodontic treatment (4 studies; average of 3.2 ΔE units; 95% CI=2.0-4.4 ΔE units), which was more than the variation among controls (1 study; MD=1.9 ΔE units; 95% CI=1.7-2.2 ΔE units). However, the quality of evidence was very low, due to the inclusion of non-randomized studies, bias, and imprecision. Re-analysis of raw study data indicated significant differences in clinically discernable treatment-induced color changes were seen between chemically- and light-cured adhesives and among the various tooth categories.

Conclusion Existing evidence of very low quality indicates that orthodontic treatment might be associated with alterations of tooth color, which are not however consistently clinically discernible. Treatment induced color alterations might be dependent on bonding material and tooth type, but evidence supporting this is weak.

Short title: Orthodontics treatment and tooth color

Key words: orthodontics, orthodontic appliances, tooth color, adverse effects, clinical trials, systematic review, meta-analysis

MANUSCRIPT

Introduction

Fixed appliances have become an integral part of contemporary orthodontic treatment, as they provide a secure hold on the teeth and enable efficient tooth movement in space. Fixed appliance treatment became more widespread with the introduction of the enamel etch technique [5] that enabled a secure fix of the orthodontic appliances bonded on the teeth and the development of more simplified appliances and techniques [29] that still provided precise biomechanical control of the teeth.

Although the ability of orthodontic appliances to therapeutically produced an acceptable outcome has been documented [33], several adverse effects to the soft and hard tissues of the oral cavity have also been reported, with the most prominent being gingival recessions [24], alveolar bone loss [4], periodontal attachment loss [34], root resorption [49], white spot lesions [15], and tooth discoloration [50].

Tooth color results from scattering of light through the tooth volume and is one of the main factors contributing to the pleasantness of the smile [26]. Tooth color seems to be mainly determined by dentin, with enamel playing a lesser role through scattering at wavelengths in the blue range [45]. Ultimately, tooth color can be influenced by a combination of intrinsic traits and extrinsic stains that may be present on the tooth surface or within the dental tissues [23], like stains promoted by smoking, drinking red wine, use of cationic agents such as chlorhexidine, or metal salts such as tin and iron [30, 48].

It has long been documented that the bonding and debonding procedures of orthodontic appliance on dental enamel can affect its color [13, 39]. This can be attributed to dissolution of enamel rods during acid etching [46], absorption of particles/substances from food, plaque, or degradation and corrosion products [11, 43], potentially discoloring [12] adhesive remnants on / in the dental tissues [37], and enamel injuries in terms of substance loss [2] or surface roughness [7, 12] during adhesive removal.

A previous systematic review on the subject reported that there is no strong evidence that orthodontic treatment with fixed appliances alters the original color of enamel [6]. However, that review included also in vitro studies, which might not reliably reflect the clinical situation [25]. Additionally, the last review covered studies published up to 2014, which means that it is possibly outdated [41]. Therefore, aim of the present systematic review was to critically assess evidence from longitudinal clinical studies on the effect of orthodontic treatment on tooth color and to answer the research question: Does orthodontic treatment have a permanent effect on tooth color of humans?

Material and methods

Protocol, eligibility criteria, and registration

The protocol for this review was made *a priori* based on the PRISMA-P statement [40], registered in PROSPERO (CRD42017064862), and all *post hoc* changes were appropriately noted. According to the PICOS schema, parallel or split-mouth randomized clinical trials and prospective non-randomized controlled or uncontrolled cohort studies on human patients assessing the effect of any kind of fixed orthodontic appliance on tooth color were included. Studies with potentially discoloring agents/treatment, studies not assessing tooth color, cross-sectional studies, animal studies, non-clinical studies, and non-relevant studies were excluded. This systematic review was conducted and reported according to Cochrane Handbook [19] and PRISMA statement [28], respectively.

Information sources and literature search

A total of nine electronic databases were searched systematically by one author (SNP) without any limitations from inception up to May 16th, 2017 (Appendix 1). Two additional sources (Google Scholar and ISRCTN registry) were manually searched for additional studies by the same author. Authors of included studies were contacted for individual patient data (Appendix 2). No limitations concerning language, publication year or status were applied. The reference lists of the included trials and relevant reviews were manually searched as well.

Study selection, data collection, and risk of bias in individual studies

Titles, abstracts, and full texts of studies identified from the literature search were screened by one author (RK) with a subsequent duplicate independent checking against the eligibility criteria by another author (SNP), while conflicts were resolved by the last author (TE). The same protocol was applied for the extraction of study characteristics (study design, setting, country, language, patient number, sex, age, malocclusion, appliances, treatment duration, oral hygiene measures, post-debond cleaning procedures, outcome measured, measurement unit, and timing) and numerical data using pre-determined and piloted extraction forms. Piloting of the forms was performed during the protocol stage until over 90% agreement was reached.

The risk of bias of the included randomized and non-randomized studies was assessed using Cochrane's risk of bias tool [19] and the ROBINS-I (Risk Of Bias In Non-randomized Studies - of Interventions) tool [44], respectively.

Data synthesis

The outcome for the meta-analysis was the overall change in tooth color (ΔE), as measured according to the CIE Lab system (Commission Internationale de l'Eclairage, L^* , a^* , b^*). The CIE colour L^* parameter corresponds to the value or degree of lightness in the Munsell system, whereas the a^* and b^* co-ordinates designate positions on red/ green ($+a^*$ = red, $-a^*$ = green) and yellow/blue ($+b^*$ = yellow, $-b^*$ = blue) axes. As the effects of orthodontic fixed appliance treatment is bound to be affected by many factors, including type of orthodontic appliances and adhesives, treatment duration, and patient-specific characteristics [3, 13, 50], a random-effects model was deemed appropriate to encompass this variability and calculate the average distribution of true effects among studies. The novel Paule-Mandel random-effects estimator was used instead of the DerSimonian and Laird one, based on recent guidance [47]. Mean Differences (MD) of treatment-induced changes between treated and untreated patients with their corresponding 95% Confidence Intervals (CI) were chosen as the primary outcome from controlled studies. Additionally, average treatment-induced changes among treated patients only were calculated as the secondary outcome for uncontrolled studies.

Absolute and relative between-trial heterogeneity was quantified with the τ^2 metric and the I^2 statistic, respectively. The latter is defined as the proportion of total variability in the results explained by heterogeneity, and not chance [17]. 95% CIs around all heterogeneity measures were calculated to quantify existing uncertainty. 95% predictive intervals were calculated for meta-analyses of three trials or more, to incorporate existing heterogeneity and provide a clinically-relevant range of possible effects for a future clinical setting [18]. Re-analysis of individual patient data provided by the authors were performed with generalized linear regression that accounted for within-person clustering of teeth and setting a change of 3.7 ΔE units as a cut-off for clinically perceivable color change [22]. All analyses were run in Stata SE 14.0 (StataCorp, College Station, TX) by one author (SNP). A two-tailed P-value of 0.05 was considered significant for hypothesis-testing, except for a 0.10 used for the test of heterogeneity and reporting biases.

Additional analyses and risk of bias across studies

Possible sources of heterogeneity were planned in the protocol to be sought through mixed-effects subgroup analyses and random-effects meta-regression for meta-analyses of \geq five studies. Eventually, subgroup analyses were only descriptively given, but not formally conducted, as less than five studies existed. The same goes for the assessment

of reporting biases and publication bias that couldn't be formally assessed as <10 studies were finally included. The overall quality of evidence (confidence in effect estimates) for each outcome was rated using the GRADE approach [16]. The minimal clinical important, large, and very large effects were to augment the produced forest plots with contours of effect magnitude [32]. Robustness of the results were planned to be checked through sensitivity analysis controlling for of bias originating from biased study designs, inconsistency, or reporting biases.

Results

Study selection

The literature search yielded a total of 427 hits (Fig. 1), 32 of which proceeded to full text assessment after eliminating duplicates and ineligible studies by title or abstract (Appendix 3). Finally, a total of four papers were identified as eligible for inclusion in the present systematic review, which pertained to four unique clinical studies published in English between 2010 and 2015 were included. Apart from data from published reports, we attempted to procure individual patient data from all included studies to reanalyze, however only one answered and provided raw data [25].

Study characteristics

The descriptive characteristics of the four included studies can be seen in Tables 1-2. From these, one was a randomized trial comparing self-etching and conventional adhesives, one was a prospective controlled cohort study comparing treated and untreated patients, one was a prospective uncontrolled cohort study comparing light-cured to chemically-cured adhesives, and the last was a prospective uncontrolled cohort study of conventional appliances. The included studies were conducted in university clinics in Greece, Iraq, Jordan, and Turkey. Overall, 138 patients were included with a mean age of 15.7 years (from the 3 studies reporting mean age) and with 46.4% patients being male (from the 4 studies reporting sex). All studies assessed color changes of anterior/posterior teeth at either tooth or patient level.

Risk of bias within studies

The risk of bias assessment is reported in Table 3 for the identified studies. Both the one randomized and all three non-randomized studies were found to be in high or serious risk of bias. The most problematic issues were confounding, lack of blinding, and selective reporting or missing data.

Results of individual studies and data synthesis

Most of the analyses were based on the data included in the published report. The only exception was the study of Karamouzou et al. [25], where the authors provided raw data for re-analysis (Appendix 4-7). The results indicated that orthodontic treatment lead to an average change by about 2.8 ΔE units (Appendix 4). Additionally, teeth bonded with chemically-cured composite were more heavily discolored than those bonded with light cured composite (difference of 0.4 ΔE units; $P < 0.001$; Appendix 5). Also, considerable variance was found among teeth, with canines being the least discolored, followed by central incisors, 1st premolars, and finally lateral incisors, which were the most discolored teeth (Appendix 5). No difference was found between upper versus lower teeth or right versus left teeth. As far as clinical relevance is concerned, 13% of all treated teeth had color changes that could be clinically discernible (change ≥ 3.7 ΔE units). Teeth bonded with chemically-cured composite were in 158% higher risk of clinically-relevant discoloration than those bonded with light-cured composite (Appendix 6). Finally, lateral incisors had 197% higher risk of clinically-relevant discoloration compared to canines (Appendix 6).

As far as analysis of the primary outcome from controlled data is concerned, evidence from one study indicated that orthodontic treatment was associated with an overall statistically significant tooth color change 4-9 months into treatment by 1.9 ΔE units (95% CI=1.7 to 2.2) compared to untreated patients (Table 4). However, the quality of evidence according to the GRADE approach was judged as very low, due to the non-randomized nature of the study, bias, and imprecision.

Meta-analysis of the secondary outcome of treatment-induced color changes of the teeth of treated patients are given in Table 5. Meta-analysis of four studies and 946 teeth indicated that the color of orthodontically treated teeth changed on average by 3.2 ΔE units (95 CI=2.0-4.4) during or after orthodontic treatment compared to the pre-treatment color. However, extreme absolute or relative heterogeneity was observed, which led to a very imprecise prediction interval of -3.2 to 10.1 ΔE units that was inconclusive.

Additional analyses

As the primary outcome from controlled data was reported only by a single non-randomized study, sensitivity analyses could be performed only for the secondary outcome of uncontrolled data. Sensitivity analyses either to alleviate the extreme heterogeneity seen or to omit the potentially biased non-randomized study designs led to similar results, indicating that orthodontic treatment was associated with an average tooth color change by about 2.8-2.9 ΔE units,

which was slightly lower than the original analysis (Table 6). Subgroup analyses were planned a priori but could not eventually be conducted formally. Evidence of different color change pattern might be seen between chemically- and light-cured resins (as stated before) and according to the vertical thirds of each crown surface (incisal, middle, and apical third), but formal analyses are needed to confirm or refute them.

Discussion

The present systematic review summarized evidence from clinical controlled and uncontrolled studies on the effect of orthodontic treatment on tooth color changes. The results of the four identified studies, including 946 teeth, indicated that orthodontic treatment with fixed appliances might have a significant effect on tooth color, although this is not always clinically discernable. After active treatment, the average tooth color becomes darker and shifts into more red and especially yellow color ranges [25, 27]. Possible explanations for these changes include enamel crystal dissipation during etching [36], resin tag penetration in enamel [13], development of white spot lesions [31], absorption of food colorants [21] or corrosion products [20], enamel morphology alteration by debonding and adhesive removal [51]. Additionally, re-analysis of raw data from a single study [25] indicated that about 13% of all bonded teeth present clinically discernable color changes, but this is more prevalent on patient level, where only 20% of patients have no clinically discernable discolorations and the rest have at least one tooth with changed color (Appendix 4).

Considerable differences were seen in the treatment-induced color change among the various bonded teeth with canines being the least affected, followed by central incisors, 1st premolars, and ultimately lateral incisors, which were most affected (Appendix 5-6) [25]. Another study reported no effect of tooth type on treatment-induced color change [1], but raw data were not available for re-analysis.

There are many factors that might possibly influence the effect of orthodontic treatment on tooth color, including among others patient characteristics, etching/bonding protocol, characteristics of the orthodontic appliance, treatment duration, post-debond cleaning protocol, oral hygiene during treatment, and the patient's dietary preferences. However, no evidence currently exists about these and well-designed clinical studies with adequate sample and appropriate analysis are needed to investigate these and minimize the adverse effects of orthodontics on tooth color.

Limited evidence indicated that chemically-cured composite resins used to bond the orthodontic appliances lead to significantly more discoloration to a clinically relevant point than light-cured composite resins (Appendix 5-6) [25]. This might be attributed to chemical differences between the two resins [10, 12, 14] like filler content and

polymerization conversion, which might affect their color stability. Another study reported no color differences between teeth bonded with self-etching or conventional etch & prime protocol [1]. This was attributed to the similarity between the two protocols of the etching pattern at the uppermost enamel layer, which discolors first, independently of penetration depth [8]. However, caution is warranted by the interpretation of these data, as evidence was of poor quality, due to the non-randomized study design and its inherent bias. Further evidence from well-designed randomized trials is needed to confirm or refute this finding.

Although the present systematic review has several strengths including a priori registration [42], extensive literature search, inclusion of individual patient data, and appropriate methods to risk of bias assessment [19, 44], data synthesis [47], and critical appraisal [16], several limitations also existed. First of all, only a handful of studies could be eventually included in the review, which affects the results' robustness and precluded the conduct of many planned analyses (Appendix 7). Additionally, three out of the four included studies were non-randomized, which means their conclusions might be potentially biased [35]. Finally, although color assessments with the CIE Lab protocol are more objective and consistent than subjective visual inspections of color, it still is prone to systematic and random errors [9, 38].

The results of the present study are applicable to adolescent and post-adolescent patients that receive fixed appliance treatment in orthodontic university clinics. The results might be less generalizable to pre-adolescent or older adult patients or patients that seek treatment in other clinical settings.

Conclusion

The present systematic review of in vivo clinical evidence indicates that fixed appliance treatment seems associated with statistically significant alteration of tooth color. These color changes are not always visible clinically and considerable difference are seen according to the bonding material and the various teeth. However, existing evidence is limited and based mainly on potentially flawed study designs and therefore additional studies are needed in order to robustly assess the effect of orthodontic treatment on tooth color and to find ways to minimize this.

References

1. Al Maaitah EF, Abu Omar AA, Al-Khateeb SN (2013) Effect of fixed orthodontic appliances bonded with different etching techniques on tooth color: A prospective clinical study. *Am J Orthod Dentofacial Orthop* 144:43–49
2. Al Shamsi AH, Cunningham JL, Lamey PJ, Lynch E (2007) Three-dimensional measurement of residual adhesive and enamel loss on teeth after debonding of orthodontic brackets: an in-vitro study. *Am J Orthod Dentofacial Orthop* 131:301.e9–15
3. Arnold S, Koletsi D, Patcas R, Eliades T (2016) The effect of bracket ligation on the periodontal status of adolescents undergoing orthodontic treatment. A systematic review and meta-analysis. *J Dent* 54:13–24
4. Bollen AM, Cunha-Cruz J, Bakko DW, Huang GJ, Hujoel PP (2008) The effects of orthodontic therapy on periodontal health: a systematic review of controlled evidence. *J Am Dent Assoc* 139:413–422
5. Buonocore MG, Matsui A, Gwinnett AJ (1968) Penetration of resin dental materials into enamel surfaces with reference to bonding. *Arch Oral Biol* 13:61–70
6. Chen Q, Zheng X, Chen W, Ni Z and Zhou Yu (2015) Influence of orthodontic treatment with fixed appliances on enamel color: a systematic review. *BMC Oral Health* 15:13
7. Chung K (1994) Effects of finishing and polishing procedures on the surface texture of resin composites. *Dent Mater* 10:325–330
8. Cinader D (2001) Chemical processes and performance comparisons of Transbond Plus self-etching primer. *Orthod Perspect* 8:5–6
9. Douglas RD (1997) Precision of in vivo colorimetric assessments of teeth. *J Prosthet Dent* 77:464–70
10. Eldiwany M, Friedl KH, Powers JM (1995) Color stability of light-cured and post-cured composites. *Am J Dent* 8:179–181
11. Eliades T, Athanasiou AE (2002) In vivo aging of orthodontic alloys: implications for corrosion potential, nickel release, and biocompatibility. *Angle Orthod* 72:222–237
12. Eliades T, Gioka C, Heim M, Eliades G, Makou M (2004) Color stability of orthodontic adhesive resins. *Angle Orthod* 74:391–393
13. Eliades T, Kakaboura A, Eliades G, Bradley TG (2001) Comparison of enamel colour changes associated with orthodontic bonding using two different adhesives. *Eur J Orthod* 23:85–90

14. Gioka C, Bourauel C, Hiskia A, Kletsas D, Eliades T, Eliades G (2005) Light-cured or chemically cured orthodontic adhesive resins? A selection based on the degree of cure, monomer leaching, and cytotoxicity. *Am J Orthod Dentofacial Orthop* 127:413–419
15. Gorelick L, Geiger AM, Gwinnett AJ (1982) Incidence of white spot formation after bonding and banding. *Am J Orthod* 81:93–98
16. Guyatt GH, Oxman AD, Schünemann HJ, Tugwell P, Knottnerus A (2011) GRADE guidelines: a new series of articles in the Journal of Clinical Epidemiology. *J Clin Epidemiol* 64:380–382
17. Higgins JP, Thompson SG, Deeks JJ, Altman DG (2003) Measuring inconsistency in meta-analyses. *BMJ* 327:557–560
18. Higgins JP, Thompson SG, Spiegelhalter DJ (2009) A re-evaluation of random-effects meta-analysis. *J R Stat Soc Ser A Stat Soc* 172:137–159
19. Higgins JPT, Green S. Cochrane handbook for systematic reviews of interventions version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. <http://www.cochrane-handbook.org>. Accessed 15 August, 2017.
20. Hodges SJ, Spencer RJ, Watkins SJ (2000) Unusual indelible enamel staining following fixed appliance treatment. *J Orthod* 27:303–6
21. Inokoshi S, Burrow MF, Kataumi M, Yamada T, Takatsu T (1996) Opacity and color changes of tooth-colored restorative materials. *Oper Dent* 21:73–80
22. Johnston WM, Kao EC (1989) Assessment of appearance match by visual observation and clinical colorimetry. *J Dent Res* 68:819–822
23. Joiner A, Hopkinson I, Deng Y, Westland S (2008) A review of tooth colour and whiteness. *J Dent* 36 Suppl 1:S2–7
24. Joss-Vassalli I, Grebenstein C, Topouzelis N, Sculean A, Katsaros C (2010) Orthodontic therapy and gingival recession: a systematic review. *Orthod Craniofac Res* 13:127–141
25. Karamouzos A, Athanasiou AE, Papadopoulos MA, Kolkithas G (2010) Tooth color assessment after orthodontic treatment: a prospective clinical trial. *Am J Orthod Dentofacial Orthop* 138:537.e1–8

26. Kershaw S, Newton JT, Williams DM (2008) The influence of tooth colour on the perceptions of personal characteristics among female dental patients: comparisons of unmodified, decayed and 'whitened' teeth. *Br Dent J* 204:E9
27. Kim SP, Hwang IN, Cho JH, Hwang HS (2006) Tooth color changes associated with the bracket bonding and debonding. *Korean J Orthod* 36:114–124
28. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JP, Clarke M, Devereaux PJ, Kleijnen J, Moher D (2009) The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *J Clin Epidemiol* 62:e1–e34
29. Mousouleas S, Papageorgiou SN, Eliades T (2017) Treatment effects of various prescriptions and techniques for fixed orthodontic appliances: A systematic review. *J Orofac Orthop*. doi: 10.1007/s00056-017-0094-0
30. Nathoo SA (1997) The chemistry and mechanisms of extrinsic and intrinsic discoloration. *J Am Dent Assoc* 128 Suppl:6S–10S
31. Øgaard B, Rolla G, Arends J (1988) Orthodontic appliances and enamel demineralization. Part 1. Lesion development. *Am J Orthod Dentofacial Orthop* 94:68–73
32. Papageorgiou SN (2014) Meta-analysis for orthodontists: Part II--Is all that glitters gold? *J Orthod* 41:327–336
33. Papageorgiou SN, Höchli D, Eliades T (2017) Outcomes of comprehensive fixed appliance orthodontic treatment: A systematic review with meta-analysis and methodological overview. *Korean J Orthod* 47:401–413
34. Papageorgiou SN, Papadelli AA, Eliades T (2017) Effect of orthodontic treatment on periodontal clinical attachment: a systematic review and meta-analysis. *Eur J Orthod*. doi: 10.1093/ejo/cjx052
35. Papageorgiou SN, Xavier GM, Cobourne MT (2015) Basic study design influences the results of orthodontic clinical investigations. *J Clin Epidemiol* 68:1512–1522
36. Retief DH, Busscher HJ, de Boer P, Jongebloed WL, Arends J (1986) A laboratory evaluation of three etching solutions. *Dent Mater* 2:202–206
37. Ribeiro AA, Almeida LF, Martins LP, Martins RP (2017) Assessing adhesive remnant removal and enamel damage with ultraviolet light: An in-vitro study. *Am J Orthod Dentofacial Orthop* 151:292–296
38. Russell MD, Gulfranz M, Moss BW (2000) In-vivo measurement of colour changes in natural teeth. *J Oral Rehabil* 27:786–792
39. Sandison RM (1981) Tooth surface appearance after debonding. *Br J Orthod* 8:199–201

40. Shamseer L, Moher D, Clarke M, Gherzi D, Liberati A, Petticrew M, Shekelle P, Stewart LA; PRISMA-P Group (2015) Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ* 349:g7647
41. Shojania KG, Sampson M, Ansari MT, Ji J, Doucette S, Moher D (2007) How quickly do systematic reviews go out of date? A survival analysis. *Ann Intern Med* 147:224–233
42. Sideri S, Papageorgiou SN, Eliades T (2017) Are orthodontic systematic reviews registered a priori in PROSPERO? *J Orthod* 7:1–7
43. Sifakakis I, Eliades T (2017) Adverse reactions to orthodontic materials. *Aust Dent J* 62 Suppl 1:20–28
44. Sterne JA, Hernán MA, Reeves BC, Savović J, Berkman ND, Viswanathan M, Henry D, Altman DG, Ansari MT, Boutron I, Carpenter JR, Chan AW, Churchill R, Deeks JJ, Hróbjartsson A, Kirkham J, Jüni P, Loke YK, Pigott TD, Ramsay CR, Regidor D, Rothstein HR, Sandhu L, Santaguida PL, Schünemann HJ, Shea B, Shrier I, Tugwell P, Turner L, Valentine JC, Waddington H, Waters E, Wells GA, Whiting PF, Higgins JP (2016) ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ* 355:i4919
45. ten Bosch JJ, Coops JC (1995) Tooth color and reflectance as related to light scattering and enamel hardness. *J Dent Res* 74:374–380
46. Tong LS, Pang MK, Mok NY, King NM, Wei SH (1993) The effects of etching, micro-abrasion, and bleaching on surface enamel. *J Dent Res* 72:67–71
47. Veroniki AA, Jackson D, Viechtbauer W, Bender R, Bowden J, Knapp G, Kuss O, Higgins JP, Langan D, Salanti G (2016) Methods to estimate the between-study variance and its uncertainty in meta-analysis. *Res Synth Methods* 7:55–79
48. Watts A, Addy M (2001) Tooth discolouration and staining: a review of the literature. *Br Dent J* 190:309–316
49. Weltman B, Vig KW, Fields HW, Shanker S, Kaizar EE (2010) Root resorption associated with orthodontic tooth movement: a systematic review. *Am J Orthod Dentofacial Orthop* 137:462–476
50. Zachrisson BJ (1977) A posttreatment evaluation of direct bonding in orthodontics. *Am J Orthod* 71:173–189.
51. Zarrinnia K, Eid NM, Kehoe MJ (1995) The effect of different debonding techniques on the enamel surface: an in vitro qualitative study. *Am J Orthod Dentofacial Orthop* 108:284–293

TABLES

Tab. 1 Patient demographics of the included studies

Study ID	Design; setting; country; language	Patients (M/F); Age in years*	Malocclusion; Tx plan	Tx duration*	Appliances; Ligs	OH status; OH measures	Post-debond cleaning
Al Maaitah 2013	RCT _{sm} ; Uni; JOR; EN	EG1/EG2: 34 (13/21); 18.6	EG1/EG2: NR; NR	12.0-15.0 mos	SS BRs; NR EG1: SEP EG2: conv primer	Good OH; OH instructions	Spiral 12-fluted tungsten carbide bur (low-speed)
Corekci 2015	pNRS _{sm/par} ; Uni; TUR; EN	EG1-4: 22 (11/11); 14.4 CG: 22 (10/12); 15.1	EG1-4: crowding<4 mm; NR CG: no crowding; NR	EG1-4: 4.3 mos CG: 8.5 mos	SS BRs; SS Ligs EG1-4: 4 different ADHs	Good OH; OH assessment/ motivation	Carbide bur (high-then low-speed); sof-lex polishing discs
Karamouzos 2010	pNRS _{sm} ; Uni; GR; EN	EG1-2: 26 (13/13); 13.6	EG1-2: crowding<4 mm NR; NR	18.0-26.0 mos	SS BRs EG1: chem-cured ADH EG2: light-cured ADH	Good OH; OH assessment and motivation at every appointment	Carbide bur (low-speed)
Al-Laban 2015	pNRS; Uni; IRQ; EN	EG: 34 (17/17); 18.0-40.0	NR; NR	NR	SS BRs	Good OH; OH instructions; bad-OH patients excluded	12 fluted tungsten carbide bur (low-speed); extra fine sof-lex polishing discs

*when one number is given, this pertains to the mean; if mean is not reported, then the range is given as two values.

BR, bracket; CG, control (untreated) group; Conv, conventional; ADH, adhesive; EG, experimental (treated) group; Ligs, ligatures; NR, not reported; OH, oral hygiene; par, parallel design; pNRS, prospective non-randomized study; RCT, randomized clinical trial; RCT_{sm}, split mouth randomized clinical trial; SEP, self-etching primer; sm, split mouth (or within-patient group allocation) design; SS, stainless steel; Tx, treatment; uni, university.

Tab. 2 Outcome characteristics of the included studies

Study	Outcome	Measured teeth	Measurement unit	Outcome timing
Al Maaitah 2013	CIE L*a*b*	Max/Mand 3-3	Tooth	<ul style="list-style-type: none">▪ Pre-bond▪ 1 wk post-debond
Corekci 2015	CIE L*a*b*	42, 41, 31, 32	Tooth	<ul style="list-style-type: none">▪ Pre-bond▪ 4.3-8.5 mos after bond
Karamouzos 2010	CIE L*a*b*	Max/Mand 4-4	Tooth	<ul style="list-style-type: none">▪ Pre-bond▪ Post-debond
Al-Laban 2015	CIE L*a*b*	3-3	Patient	<ul style="list-style-type: none">▪ Pre-bond▪ Post-debond

CIE L*a*b*, color space as specified by the International Commission on Illumination; Mand, mandibular; Max, maxillary; mo, month; wk, week.

Tab. 3 Risk of bias assessment of randomized and non-randomized identified studies

Bias domain for RCTs	Al-Maaitah 2013		Bias domain for non-RCTs	Al Laban 2015	Corekci 2015	Karamouzos 2010
Sequence generation	Unclear		Confounding	Serious	Serious	Moderate
Allocation concealment	Low		Selection of participants	Low	Low	Moderate
Blinding of participants/ personnel	Unclear		Classification of interventions	Low	Low	Low
Blinding of outcome assessors	High		Deviations from intended interventions	Low	Low	Low
Incomplete outcome data	High		Missing data	Serious	Serious	Serious
Selective outcome reporting	High		Measurement of outcomes	Moderate	Low	Moderate
Other sources of bias	Unclear		Selection of reported results	Moderate	Moderate	Moderate
Overall	High		Overall	Serious	Serious	Serious

RCT, randomized clinical trial.

Tab. 4 Results of the single identified controlled clinical study comparing treated and untreated patients

Study (teeth)	Comparison	Timing	Outcome	Mean Difference	95% Confidence Interval	P
1 (176)	Treatment vs Control	4.3-8.5 months after bonding	ΔE	1.94	1.71,2.18	<0.001

Tab. 5 Summary of findings table according to the GRADE approach based on the single identified controlled clinical study comparing treated and untreated patients

Outcome Trials (patients) – follow-up	Anticipated absolute effects			Quality of the evidence (GRADE) ^a	What happens to treated teeth
	Control*	Tx	Difference Tx-control (95% CI)		
Change in <i>E</i> (total color difference) 1 trial (176 teeth) – 4-9 mos	ΔE is increased by 0.29	-	1.94 increase (1.71 to 2.18 increase)	⊕○○○ very low ^{a,b} due to bias, imprecision	May lead to increase in total color difference ΔE .

Effects of fixed appliance treatment on tooth color.

Patient or population: pre- or post-adolescence non-adult patients attending an orthodontic university clinic.

Settings: university clinics (Turkey).

* Reponse in the untreated group of the one included controlled clinical study.

^a Starts from "low", due to the inclusion of non-randomized studies. Downgraded further by one point due to serious limitations (high risk of bias).

^b Downgraded by one due to imprecision originating from the inclusion of a small sample size.

Abbreviations

CI, confidence interval; GRADE, Grading of Recommendations Assessment, Development and Evaluation; Tx, treatment.

Tab. 6 Meta-analysis of all 4 identified controlled or uncontrolled clinical studies on tooth color changes (before & after treatment) of treated patient

Outcome	Studies (teeth)	Tx change	95% CI	P	I ² (95% CI)	t ² (95% CI)	95% prediction
<i>Original analysis</i>							
<i>ΔE</i>	4 (946)	3.21	2.03,4.38	<0.001	100% (100%-100%)	1.42 (0.44-20.0)	-3.63,10.05
<i>Sensitivity analysis (Two studies omitted to alleviate extreme heterogeneity)</i>							
<i>ΔE</i>	2* (824)	2.84	2.80,2.88	<0.001	32% (0-100%)	0 (0-1.27)	NA
<i>Sensitivity analysis (three non-randomized studies omitted)</i>							
<i>ΔE</i>	1 (408)	2.85	2.83,2.87	<0.001	NA	NA	NA

CI, confidence interval; NA, non-applicable; Tx, treatment.


Tab. 7 Explorative assessment of subgroup differences in total color change ΔE according to identified controlled or uncontrolled clinical studies

Studies (teeth)	Subgroup	Tx Change	95% CI	P _{SG}
1 (208)	Maxillary teeth	2.75	2.63,2.87	NA
1 (208)	Mandibular teeth	2.86	2.74,2.98	
1 (17)	Female patients	4.30	3.77,4.83	NA
1 (17)	Male patients	4.14	3.69,4.59	
1 (208)	Anterior teeth (incisors)	2.82	2.70,2.94	NA
1 (208)	Posterior teeth (canine and premolar)	2.80	2.70,2.90	
1 (208)	Light-cured adhesive	2.60	2.50,2.70	NA
1 (208)	Chemically-cured adhesive	3.00	2.88,3.12	
1 (17)	Incisal tooth third	3.64	3.29,3.99	NA
1 (17)	Middle tooth third	4.96	4.65,5.27	
1 (17)	Gingival tooth third	4.05	3.68,4.42	

CI, confidence interval; NA, non-applicable; P_{SG}, p value for subgroup differences; Tx, treatment.

Figure Captions

Fig. 1. PRISMA flow diagram for the identification and selection of studies for this systematic review



Does orthodontic treatment have a permanent effect on tooth color? A systematic review and meta-analysis

Appendix 1. Search strategy for each database with the corresponding hits (last search 16.5.2017)

Nr	Database	Search	Limit	Hits
1	PubMed	(orthodon* OR malocclusion) AND (treatment OR therapy OR brace* OR bracket* OR "fixed appliance") AND (tooth OR teeth) AND (color* OR colour* OR discolor* OR discolour*)	(Human)(Clinical Study; Clinical Trial; Clinical Trial, Phase I-IV; Comparative Study; Controlled Clinical Trial; Multicenter Study; Observational Study; Pragmatic Clinical Trial; Randomized Controlled Trial)	110
2	CDSR	Same	-	0
3	CENTRAL	Same	-	37
4	DARE	Same	-	0
5	Embase	Same	-	76
6	VHL	Same	Humans	47
7	Scopus	Same	(Dentistry)(Human; Humans)(Controlled Study; Randomized Controlled Trial; Controlled Clinical Trial; Clinical Trial)	42
8	WoK	(orthodon* OR malocclusion) AND (treatment OR therapy OR brace* OR bracket* OR "fixed appliance") AND (tooth OR teeth) AND (color* OR colour* OR discolor* OR discolour*)	DENTISTRY ORAL SURGERY MEDICINE	110
9	ClinicalTrials	orthodontic AND (color OR colour OR discoloration OR discolouration)		3
		SUM (with overlap)		425
		SUM (without overlap)		297

Appendix 2. Authors contacted for individual patient data.

Nr	Paper	
1	Al Maaith EF, Abu Omar AA, Al-Khateeb SN. Effect of fixed orthodontic appliances bonded with different etching techniques on tooth color: a prospective clinical study. Am J Orthod Dentofacial Orthop. 2013;144(1):43-9.	E-mailed October 23rd; awaiting response.
2	Al-Laban YRA. Comparison of enamel color alteration between bonded and free unbonded surfaces of maxillary anterior teeth after fixed orthodontic therapy : A Prospective Clinical Study. J Baghdad College Dentistry 2015;27:174-178.	E-mail could not be found
3	Corekci B, Toy E, Ozturk F, Malkoc S, Ozturk B. Effects of contemporary orthodontic composites on tooth color following short-term fixed orthodontic treatment: a controlled clinical study. Turk J Med Sci. 2015;45(6):1421-8.	E-mailed October 23rd; awaiting response.
4	Karamouzou A, Athanasiou AE, Papadopoulos MA, Kolokithas G. Tooth-color assessment after orthodontic treatment: a prospective clinical trial. Am J Orthod Dentofacial Orthop. 2010;138(5):537.e1-8; discussion -9.	E-mailed October 23rd; responded and provided individual patient raw data.

Appendix 3. List of all identified studies together with their selection status.

Nr	Paper	Status
1	Agrawal A, Shigli A. Comparison of six different methods of cleaning and preparing occlusal fissure surface before placement of pit and fissure sealant: an in vitro study. <i>J Ind Soc Pedodont Prev Dent</i> 2012;30(1):51-5.	Exclusion by title
2	Ahrari F, Akbari M, Mohammadpour S, Forghani M. The efficacy of laser-assisted in-office bleaching and home bleaching on sound and demineralized enamel. <i>Laser Therapy</i> . 2015;24(4):257-64.	Exclusion by title
3	Akin M, Baka ZM, Ileri Z, Basciftci FA. Can demineralized enamel surfaces be bonded safely? <i>Acta Odontologica Scandinavica</i> . 2014;72(4):283-9.	Exclusion by title
4	Akyalcin S, Alexander SP, Silva RM, English JD. Evaluation of three-dimensional root surface changes and resorption following rapid maxillary expansion: a cone beam computed tomography investigation. <i>Orthod Craniofac Res</i> . 2015;18 Suppl 1:117-26.	Exclusion by title
5	Albino LGB, Gradinar O, Bertholdo G, Cruz M. Otimizeando a composição do sorriso com resina composta direta. <i>Clín int j braz dent</i> . 2013;9(3):262-70.	Exclusion by title
6	Alkilzy M, Berndt C, Meller C, Schidlowski M, Splieth C. Sealing of proximal surfaces with polyurethane tape: a two-year clinical and radiographic feasibility study. <i>J Adhes Dent</i> . 2009;11(2):91-4.	Exclusion by title
7	Almeida FT, Leite AF, De Souza Figueiredo PT, Melo NS, Sousa JB, Almeida R, et al. Dento-osseous anomalies associated to familial adenomatous polyposis mimicking florid cemento-osseous dysplasia. <i>Journal of Cranio-Maxillofacial Surgery</i> . 2012;40(8):e498-e502.	Exclusion by title
8	Almeida RC, Cevdanes LHS, Carvalho FAR, Motta AT, Almeida MAO, Styner M, et al. Soft tissue response to mandibular advancement using 3D CBCT scanning. <i>International Journal of Oral and Maxillofacial Surgery</i> . 2011;40(4):353-9.	Exclusion by title
9	Aloysius AP, Vijayalakshmi D, Deepika, Soundararajan NK, Manohar VN, Khan N. Comparative evaluation of frictional properties, load deflection rate and surface characteristics of different coloured TMA archwires - An invitro study. <i>Journal of Clinical and Diagnostic Research</i> . 2015;9(12):ZC26-ZC9.	Exclusion by title
10	Alves dos Santos MP, Luiz RR, Maia LC. Randomised trial of resin-based restorations in Class I and Class II beveled preparations in primary molars: 48-month results. <i>J Dent</i> . 2010;38(6):451-9.	Exclusion by title
11	Andersson L, Emami-Kristiansen Z, Hogstrom J. Single-tooth implant treatment in the anterior region of the maxilla for treatment of tooth loss after trauma: a retrospective clinical and interview study. <i>Dental Traumatology</i> . 2003;19(3):126-31.	Exclusion by title
12	Andrade MR, Salazar SL, de Sa LF, Portela M, Ferreira-Pereira A, Soares RM, et al. Role of saliva in the caries experience and calculus formation of young patients undergoing hemodialysis. <i>Clin Oral Invest</i> . 2015;19(8):1973-80.	Exclusion by title
13	Argueta-Figueroa L, Scougall-Vilchis RJ, Morales-Luckie RA, Olea-Mejia OF. An evaluation of the antibacterial properties and shear bond strength of copper nanoparticles as a nanofiller in orthodontic adhesive. <i>Aust Orthod J</i> . 2015;31(1):42-8.	Exclusion by title
14	Artun J, Grobety D. Periodontal status of mandibular incisors after pronounced orthodontic advancement during adolescence: A follow-up evaluation. <i>Am J Orthod Dentofacial Orthop</i> 2001;119(1):2-10.	Exclusion by title
15	Associação Brasileira de Cirurgias-DentistasPedraza H, s.afSantos CRd, Associação Odontológica de Ribeirão PretoTakeuchi CYG, Universidade Estadual PaulistaAndrade MFd. Reabilitação estética do sorriso por meio do sistema cerâmico dissilicato de lítio – relato de caso. <i>Full dent sci</i> . 2015;6(21):112-7.	Exclusion by title
16	Bacci C, Lucchiari N, Valente M, Della Barbera M, Frigo AC, Berengo M. Intra-oral bone harvesting: two methods compared using histological and histomorphometric assessments. <i>Clin Oral Implants Res</i> . 2011;22(6):600-5.	Exclusion by title
17	Baek J, Wilder-Smith P, Na J, Lee BH. Periodontal tissue changes can be imaged during orthodontic tooth movement. <i>Lasers in Surgery and Medicine</i> . 2010;42:49-50.	Exclusion by title
18	Bakor SF, Pereira JCM, Frascino S, Ladalardo TCCGP, Pignatari SSN, Weckx LLM. Desmineralização dentária de pacientes respiradores orais submetidos à expansão maxilar. <i>Braz J Otorrinolaryngol</i> . 2010;76(6):709-12.	Exclusion by title
19	Ballard RW, Hagan JL, Phaup AN, Sarkar N, Townsend JA, Armbruster PC. Evaluation of 3 commercially available materials for resolution of white spot lesions. <i>Am J Orthod Dentofacial Orthop</i> . 2013;143(4 Suppl):S78-84.	Exclusion by title
20	Bartoňová M, Dostálová T, Peterka M, Kozák J, Müllerová Ž. Long-term stability of prosthetic treatment of oronasal and oroantral communications. <i>Acta Chirurgiae Plasticae</i> . 2005;47(3):85-91.	Exclusion by title
21	Batista EL, Jr., Batista FC, Novaes AB, Jr. Management of soft tissue ridge deformities with acellular dermal matrix. <i>Clinical approach and outcome after 6 months of treatment</i> . <i>J Periodontol</i> . 2001;72(2):265-73.	Exclusion by title
22	Batson ER, Cooper LF, Duqum I, Mendonça G. Clinical outcomes of three different crown systems with CAD/CAM technology. <i>Journal of Prosthetic Dentistry</i> . 2014;112(4):770-7.	Exclusion by title
23	Baty DL, Volz JE, von Fraunhofer JA. Force delivery properties of colored elastomeric modules. <i>Am J Orthod Dentofacial Orthop</i> . 1994;106(1):40-6.	Exclusion by title
24	Bauss O, Schafer W, Sadat-Khonsari R, Knosel M. Influence of orthodontic extrusion on pulpal vitality of traumatized maxillary incisors. <i>J Endod</i> . 2010;36(2):203-7.	Exclusion by title
25	Bazargani F, Jacobson S, Lennartsson B. A comparative evaluation of lingual retainer failure bonded with or without liquid resin: A randomized clinical study with 2-year follow-up. <i>Angle Orthodontist</i> . 2012;82(1):84-7.	Exclusion by title
26	Becker A, Chaushu S. Long-term follow-up of severely resorbed maxillary incisors after resolution of an etiologically associated impacted canine. <i>Am J Orthod Dentofacial Orthop</i> 2005;127(6):650-4.	Exclusion by title
27	Benson PE, Shah AA, Campbell IF. Fluoridated elastomers: effect on disclosed plaque. <i>J Orthod</i> . 2004;31(1):41-6.	Exclusion by title
28	Bharti R, Chandra A, Tikku AP, Prasad V, Shakya VK, Singhal R. Management of mucosal fenestration with external root resorption by multidisciplinary approach. <i>BMJ Case Reports</i> . 2014;2014.	Exclusion by title
29	Bidra AS, Uribe F. Successful Bleaching of Teeth with Dentinogenesis Imperfecta Discoloration: A Case Report. <i>Journal of Esthetic and Restorative Dentistry</i> . 2011;23(1):3-10.	Exclusion by title
30	Bjerklin K, Guitirokh CH. Maxillary incisor root resorption induced by ectopic canines. <i>Angle Orthod</i> . 2011;81(5):800-6.	Exclusion by title
31	Blasco Sansano R, Castellar Ponce MD, Llorca Salort N, Valero Rosique J, García Espinosa S. Estudio sobre los factores de riesgo de caries y evaluación de un test indicador del pH y revelado de la placa y la capacidad tampón de la saliva. <i>Pediatr aten prim</i> . 2009;11(41):33-47.	Exclusion by title
32	Boeira GF, Salas MMS, Araújo DC, Masotti AS, Correa MB, Demarco FF. Factors influencing dental appearance satisfaction in adolescents: a cross-sectional study conducted in Southern Brazil. <i>Braz j oral sci</i> . 2016;15(1):8-15.	Exclusion by title
33	Borges AB, Caneppele TMF, Masterson D, Maia LC. Is resin infiltration an effective esthetic treatment for enamel development defects and white spot lesions? A systematic review. <i>J Dent</i> 2017;56:11-8.	Exclusion by title
34	Borges CM, Peres MA, Peres KG. Associação entre presença de oclusopatias e insatisfação com a aparência dos dentes e gengivas: estudo com adolescentes brasileiros. <i>Rev Bras Epidemiol</i> . 2010;13(4):713-23.	Exclusion by title
35	Borzabadi-Farahani A, Borzabadi-Farahani A. The association between orthodontic treatment need and maxillary incisor trauma, a retrospective clinical study. <i>Oral Surg Oral Med Oral Pathol Oral Radiol Endod</i> . 2011;112(6):e75-80.	Exclusion by title

36	Bowen RL, Rupp NW, Eichmiller FC, Stanley HR. Clinical biocompatibility of an experimental dentine-enamel adhesive for composites. <i>Int Dent J</i> . 1989;39(4):247-52.	Exclusion by title
37	Boyd RL, Hollander BN, Eakle WS. Comparison of a subgingivally placed cannula oral irrigator tip with a supragingivally placed standard irrigator tip. <i>J Clin Periodontol</i> . 1992;19(5):340-4.	Exclusion by title
38	Boyd RL. Long-term evaluation of a SnF2 gel for control of gingivitis and decalcification in adolescent orthodontic patients. <i>Int Dent J</i> . 1994;44(1 Suppl 1):119-30.	Exclusion by title
39	Brauchli LM, Senn C, Ball J, Wichelhaus A. Force levels of 23 nickel-titanium open-coil springs in compression testing. <i>Am J Orthod Dentofacial Orthop</i> 2011;139(5):601-5.	Exclusion by title
40	Brazilian Dental J, Araujo JFd, Braga EMF, Loretto SC, Souza PdARSe, Barros TAdF, et al. One-Year Evaluation of a Simplified Ethanol-Wet Bonding Technique: A Randomized Clinical Trial. <i>Braz Dent J</i> . 2013;24(3):267-72.	Exclusion by title
41	Brightman LJ, Terezhalmay GT, Greenwell H, Jacobs M, Enlow DH. The effects of a 0.12% chlorhexidine gluconate mouthrinse on orthodontic patients aged 11 through 17 with established gingivitis. <i>Am J Orthod Dentofacial Orthop</i> . 1991;100(4):324-9.	Exclusion by title
42	Brough E, Donaldson AN, Naini FB. Canine substitution for missing maxillary lateral incisors: the influence of canine morphology, size, and shade on perceptions of smile attractiveness. <i>Am J Orthod Dentofacial Orthop</i> . 2010;138(6):705.e1-9.	Exclusion by title
43	Buckley JG, Jones ML, Hill M, Sugar AW. An evaluation of the changes in maxillary pulpal blood flow associated with orthognathic surgery. <i>Br J Orthod</i> . 1999;26(1):39-45.	Exclusion by title
44	Carrillo Baracaldo JS, Álvarez Quesada C, Calatayud Sierra J. Trabajo en equipo en Odontología: la comunicación con el laboratorio dental como clave de éxito clínico. A propósito de un caso clínico de tratamiento multidisciplinar. <i>Cient dent (Ed impr)</i> . 2006;3(2):129-36.	Exclusion by title
45	Carrillo Baracaldo JS, Álvarez Quesada MdC, Costa Ferrer F. Utilización de nuevos equipamientos en Odontología. A propósito de un caso clínico de gingivectomía con láser de diodo. <i>Cient dent (Ed impr)</i> . 2004;1(1):39-43.	Exclusion by title
46	Chalub LLFH, Martins CC, Ferreira RC, Vargas AMD. Functional dentition in Brazilian adults: An investigation of social determinants of health (SDH) using a multilevel approach. <i>PLoS ONE</i> . 2016;11(2).	Exclusion by title
47	Chan KH, Fried D. Selective Removal of Dental Composite using a Rapidly Scanned Carbon Dioxide Laser. <i>Lasers in Dentistry Xvii. Proceedings of SPIE</i> . 78842011.	Exclusion by title
48	Chan KH, Hirasuna K, Fried D. Rapid and selective removal of composite from tooth surfaces with a 9.3µm CO2 laser using spectral feedback. <i>Lasers in Surgery and Medicine</i> . 2011;43(8):824-32.	Exclusion by title
49	Chashu S, Dykstein N, Ben-Bassat Y, Becker A. Periodontal status of impacted maxillary incisors uncovered by 2 different surgical techniques. <i>J Oral Maxillofac Surg</i> . 2009;67(1):120-4.	Exclusion by title
50	Chen P, Yu S, Zhu G. The psychosocial impacts of implantation on the dental aesthetics of missing anterior teeth patients. <i>British Dental Journal</i> . 2012;213(11).	Exclusion by title
51	Cholakova R, Chenchev I, Jordanova S, Oncheva D, Chenchev L. A case of compound maxillary odontoma and mandibular hypodontia. <i>Journal of IMAB - Annual Proceeding (Scientific Papers)</i> . 2016;22(3):1217-20.	Exclusion by title
52	Chu CH, Zhang CF, Jin LJ. Treating a maxillary midline diastema in adult patients A general dentist's perspective. <i>Journal of the American Dental Association</i> . 2011;142(11):1258-64.	Exclusion by title
53	Chu FCS. Clinical considerations in managing severe tooth discoloration with porcelain veneers. <i>Journal of the American Dental Association</i> . 2009;140(4):442-6.	Exclusion by title
54	Clocheret K, Willems G, Carels C, Celis JP. Dynamic frictional behaviour of orthodontic archwires and brackets. <i>Eur J Orthod</i> 2004;26(2):163-70.	Exclusion by title
55	Contreras-Bulnes R, Scougall-Vilchis RJ, Rodríguez-Vilchis LE, Centeno-Pedraza C, Olea-Mejía OF, Alcántara-Galena MDCZ. Evaluation of self-etching adhesive and Er:YAG laser conditioning on the shear bond strength of orthodontic brackets. <i>The Scientific World Journal</i> . 2013;2013.	Exclusion by title
56	Cosyn J, Verelst K. An efficacy and safety analysis of a chlorhexidine chewing gum in young orthodontic patients. <i>J Clin Periodontol</i> . 2006;33(12):894-9.	Exclusion by title
57	Czochrowska EM, Stenvik A, Zachrisson BU. The esthetic outcome of autotransplanted premolars replacing maxillary incisors. <i>Dental Traumatology</i> . 2002;18(5):237-45.	Exclusion by title
58	Dai F, Li Y, Chen G, Chen S, Xu T. A novel method for prediction of dynamic smiling expressions after orthodontic treatment: a case report. <i>Computer Methods in Biomechanics and Biomedical Engineering</i> . 2016;19(3):340-6.	Exclusion by title
59	Davari A, Yassaei S, Karandish M, Zarghami F. In vitro evaluation of microleakage under ceramic and metal brackets bonded with LED and plasma arc curing. <i>J Contemp Dent Pract</i> . 2012;13(5):644-9.	Exclusion by title
60	de Carvalho V, de Carvalho C, de Carvalho A, Borges TGF, de Carvalho V, Santiago EFA. Statistical analysis of teeth autotransplantation in Portugal's region of Chaves. <i>Acta Odontologica Scandinavica</i> . 2014;72(3):179-86.	Exclusion by title
61	De Cicco V. Cerebro-afferent vessel and pupillary basal diameter variation induced by stomatognathic trigeminal proprioception: A case report. <i>Journal of Medical Case Reports</i> . 2012;6.	Exclusion by title
62	Demirci M, Tuncer S, Oztas E, Tekce N, Uysal O. A 4-year clinical evaluation of direct composite build-ups for space closure after orthodontic treatment. <i>Clinical Oral Investigations</i> . 2015;19(9):2187-99.	Exclusion by title
63	Diniz MB, Aranha AMF, Giro EMA. Reabilitação de dentes anteriores traumatizados pela técnica da colagem de fragmentos. <i>J Health Sci Inst</i> . 2008;26(3):366-71.	Exclusion by title
64	Domínguez-Mompell Micó R, De la Cruz Fernández C, Marcianes Moreno M, Morón Duelo R, García-Camba Varela P, Varela Morales M. Incisivos en pala: frecuencia en pacientes ortodóncicos de diversas etnias. <i>Cient dent (Ed impr)</i> . 2014;11(1):15-20.	Exclusion by title
65	Dostálová T, Tauferová E, Teuberová Z, Seydlová M, Smutný V, Racek J, et al. Shape and size of dental arch: A five-year prospective study. <i>Methods of Information in Medicine</i> . 2006;45(2):191-4.	Exclusion by title
66	Dua D, Dua A. Reconstruction and intentional replantation of a maxillary central incisor with a complete vertical root fracture: A rare case report with three years follow up. <i>Journal of Clinical and Diagnostic Research</i> . 2015;9(9):ZD06-ZD9.	Exclusion by title
67	Dueled E, Goffredsen K, Trab Damsgaard M, Hede B. Professional and patient-based evaluation of oral rehabilitation in patients with tooth agenesis. <i>Clin Oral Implants Res</i> . 2009;20(7):729-36.	Exclusion by title
68	Duque C, Negrini Tde C, Sacono NT, Spolidorio DM, de Souza Costa CA, Hebling J. Clinical and microbiological performance of resin-modified glass-ionomer liners after incomplete dentine caries removal. <i>Clin Oral Investig</i> . 2009;13(4):465-71.	Exclusion by title
69	Dursun E, Savard E, Vargas C, Loison-Robert L, Cherifi H, Bdeoui F, et al. Management of Amelogenesis Imperfecta: A 15-Year Case History of Two Siblings. <i>Operative Dentistry</i> . 2016;41(6):567-77.	Exclusion by title
70	Fanghânel J. Biomechanics and biomaterials in oral rehabilitation and dental treatment. <i>Biomedizinische Technik</i> . 2008;53(5):213-4.	Exclusion by title
71	Farrow ML, Newman SM, Oesterle LJ, Shellhart WC. Filled and unfilled restorative materials to reduce enamel decalcification during fixed-appliance orthodontic treatment. <i>Am J Orthod Dentofacial Orthop</i> . 2007;132(5):578.e1-6.	Exclusion by title

72	Fernandes LMPdSR, Rodriguez MHH, Lascala CÂ. Dentinogênese imperfeita familiar: relato de caso. Rev odonto ciênc. 2008;23(2).	Exclusion by title
73	Fidalgo TKD, Pithon MM, do Santos RL, de Alencar NA, Abrahao AC, Maia LC. Influence of topical fluoride application on mechanical properties of orthodontic bonding materials under pH cycling. Angle Orthodontist. 2012;82(6):1071-7.	Exclusion by title
74	Flugge TV, Schlager S, Nelson K, Nahles S, Metzger MC. Precision of intraoral digital dental impressions with iTero and extraoral digitization with the iTero and a model scanner. Am J Orthod Dentofacial Orthop. 2013;144(3):471-8.	Exclusion by title
75	Foersch M, Schuster C, Rahimi RK, Wehrbein H, Jacobs C. A new flash-free orthodontic adhesive system: A first clinical and stereomicroscopic study. Angle Orthodontist. 2016;86(2):260-4.	Exclusion by title
76	Freiman A, Borsuk D, Barankin B, Sperber GH, Krafchik B. Dental manifestations of dermatologic conditions. Journal of the American Academy of Dermatology. 2009;60(2):289-98.	Exclusion by title
77	Fristad I, Hagen SO, Bardsen A, Molven O. Lateral breakdown of nonendodontic origin adjacent to maxillary left incisors. International Endodontic Journal. 2000;33(5):471-4.	Exclusion by title
78	Fulton AJ, Fiani N, Verstraete FJM. Canine pediatric dentistry. Veterinary Clinics of North America - Small Animal Practice. 2014;44(2):303-24.	Exclusion by title
79	Ganesh M, Shobha T. Comparative evaluation of the marginal sealing ability of Fuji VII and Concise as pit and fissure sealants. J Contemp Dent Pract. 2007;8(4):10-8.	Exclusion by title
80	Gehrke P, Lobert M, Dhom G. Reproducibility of the pink esthetic score--rating soft tissue esthetics around single-implant restorations with regard to dental observer specialization. J Esthet Restor Dent. 2008;20(6):375-84.	Exclusion by title
81	Georgescu CE, Rusu MC, Sandulescu M, Enache AM, Didilescu AC. Quantitative and qualitative bone analysis in the maxillary lateral region. Surgical and Radiologic Anatomy. 2012;34(6):551-8.	Exclusion by title
82	Ghaffar F, Fida M. Effect of extraction of first four premolars on smile aesthetics. Eur J Orthod. 2011;33(6):679-83.	Exclusion by title
83	Goncalves AF, de Oliveira Rocha R, Oliveira MD, Rodrigues CR. Clinical effectiveness of toothbrushes and toothbrushing methods of plaque removal on partially erupted occlusal surfaces. Oral Health Prev Dent. 2007;5(1):33-7.	Exclusion by title
84	Gontijo IT, Navarro RS, Ciamponi AL, Miyakawa W, Zzell DM. Color and surface temperature variation during bleaching in human devitalized primary teeth: an in vitro study. J Dent Child (Chic). 2008;75(3):229-34.	Exclusion by title
85	Gontijo IT, Navarro RS, Ciamponi AL, Zzell DM. Whitening techniques using the diode laser and halogen lamp in human devitalized primary teeth. J Dent Child (Chic). 2008;75(2):164-7.	Exclusion by title
86	Gonzalez OL, Vera J, Orbzco MS, Mancera JT, Gonzalez KV, Malagon GV. Transient Apical Breakdown and Its Relationship with Orthodontic Forces: A Case Report. Journal of Endodontics. 2014;40(8):1265-7.	Exclusion by title
87	Gordon CT, Weaver KN, Zechi-Ceide RM, Madsen EC, Tavares ALP, Oufadem M, et al. Mutations in the endothelin receptor type a cause mandibulofacial dysostosis with alopecia. American Journal of Human Genetics. 2015;96(4):519-31.	Exclusion by title
88	Griffiths GS, Moulson AM, Petrie A, James IT. Evaluation of osteocalcin and pyridinium crosslinks of bone collagen as markers of bone turnover in gingival crevicular fluid during different stages of orthodontic treatment. Journal of Clinical Periodontology. 1998;25(6):492-8.	Exclusion by title
89	Gupta A, Tavane P, Gupta PK, Tejolatha B, Lakhani AA, Tiwari R, et al. Evaluation of microleakage with total etch, self etch and universal adhesive systems in class V restorations: An in vitro study. Journal of Clinical and Diagnostic Research. 2017;11(4):ZC53-ZC6.	Exclusion by title
90	Gupta D, Nayan S, Tippanawar H, Patil G, Jain A, Momin R, et al. Are herbal mouthwash efficacious over chlorhexidine on the dental plaque? Pharmacognosy Research. 2015;7(3):277-81.	Exclusion by title
91	Haavikko K, Rantanen - Dept. dept. Pedodont. Orthodont L, Inst D, Univ H. A follow up study of injuries to permanent and primary teeth in children. Proceedings of the Finnish Dental Society. 1976;72(5):152-62.	Exclusion by title
92	Hakki SS, Korkusuz P, Berk G, Dundar N, Saglam M, Bozkurt B, et al. Comparison of Er,Cr:YSGG laser and hand instrumentation on the attachment of periodontal ligament fibroblasts to periodontally diseased root surfaces: an in vitro study. J Periodontol. 2010;81(8):1216-25.	Exclusion by title
93	Harada K, Sato M, Omura K. Blood-flow change and recovery of sensibility in the maxillary dental pulp during and after maxillary distraction: A pilot study. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontology. 2004;98(5):528-32.	Exclusion by title
94	Hartmann A, Seiler M, Oeschey P, Dreyling M, Heinig N, Beuer F. Implant placement in the esthetic zone associated with augmentation procedures. Implantologie. 2013;21(3):321-8.	Exclusion by title
95	Hemmings K, Griffiths B, Hobkirk J, Scully C. Improving occlusion and orofacial aesthetics: Tooth repair and replacement. British Medical Journal. 2000;321(7258):438-41.	Exclusion by title
96	Hino CT, Cevidanes LH, Nguyen TT, De Clerck HJ, Franchi L, McNamara JA, Jr. Three-dimensional analysis of maxillary changes associated with facemask and rapid maxillary expansion compared with bone anchored maxillary protraction. Am J Orthod Dentofacial Orthop. 2013;144(5):705-14.	Exclusion by title
97	Ho CSF, Ming Y, Foong KWC, Rosa V, Thuyen T, Seneviratne CJ. Streptococcus mutans forms xylitol-resistant biofilm on excess adhesive flash in novel ex-vivo orthodontic bracket model. Am J Orthod Dentofacial Orthop 2017;151(4):669-77.	Exclusion by title
98	Huja SS, Mason A, Fenell CE, Mo X, Hueni S, D'Atri AM, et al. Effects of short-term zoledronic acid treatment on bone remodeling and healing at surgical sites in the maxilla and mandible of aged dogs. Journal of Oral and Maxillofacial Surgery. 2011;69(2):418-27.	Exclusion by title
99	Husmann P, Bourauel C, Wessinger M, Jäger A. The frictional behavior of coated guiding archwires. J Orofac Orthop 2002;63(3):199-211.	Exclusion by title
100	Hvaring CL, Ogaard B, Birkeland K. Tooth replacements in young adults with severe hypodontia: Orthodontic space closure, dental implants, and tooth-supported fixed dental prostheses. A follow-up study. Am J Orthod Dentofacial Orthop 2016;150(4):620-6.	Exclusion by title
101	Issar R, Mazumdar D, Ranjan S, Krishna NK, Kole R, Singh P, et al. Comparative evaluation of the etching pattern of Er,Cr:YSGG & acid etching on extracted human teeth-an ESEM analysis. Journal of Clinical and Diagnostic Research. 2016;10(5):ZC01-ZC5.	Exclusion by title
102	Ittipuriphat I, Leevailoj C. Anterior Space Management: Interdisciplinary Concepts. Journal of Esthetic and Restorative Dentistry. 2013;25(1):16-30.	Exclusion by title
103	Iwase M, Ohashi M, Tachibana H, Toyoshima T, Nagumo M. Bite force, occlusal contact area and masticatory efficiency before and after orthognathic surgical correction of mandibular prognathism. International Journal of Oral and Maxillofacial Surgery. 2006;35(12):1102-7.	Exclusion by title
104	Jacobovitz M, Lima RKD. The use of calcium hydroxide and mineral trioxide aggregate on apexification of a replanted tooth: a case report. Dental Traumatology. 2009;25(3):E32-E6.	Exclusion by title

105	Jacobs C, Grimm S, Ziebart T, Walter C, Wehrbein H. Osteogenic differentiation of periodontal fibroblasts is dependent on the strength of mechanical strain. Arch Oral Biol. 2013;58(7):896-904.	Exclusion by title
106	Jia S, Wang DF. The clinical application of the everStick high-strength fiber composite resin. Chinese Journal of Tissue Engineering Research. 2014;18(3):458-63.	Exclusion by title
107	Johnston WM. Review of Translucency Determinations and Applications to Dental Materials. Journal of Esthetic and Restorative Dentistry. 2014;26(4):217-23.	Exclusion by title
108	Josefsson E, Bjerklin K, Lindsten R. Self-perceived orthodontic treatment need and prevalence of malocclusion in 18- and 19-year-olds in Sweden with different geographic origin. Swed Dent J. 2010;34(2):95-106.	Exclusion by title
109	Kallu R, Vinckier F, Politis C, Mwalili S, Willems G. Tooth transplantations: A descriptive retrospective study. International Journal of Oral and Maxillofacial Surgery. 2006;34(7):745-55.	Exclusion by title
110	Katona TR, Winkler MM, Huang J. Stress analysis of a bulk-filled class-V chemical-cured dental composite restoration. Journal of Biomedical Materials Research. 1996;31(4):445-9.	Exclusion by title
111	Kau CH. Creation of the Virtual Patient for the Study of Facial Morphology. Facial Plastic Surgery Clinics of North America. 2011;19(4):615-22.	Exclusion by title
112	Keeling SD, King GJ, McCoy EA, Valdez M. Serum and Alveolar Bone Phosphatase Changes Reflect Bone Turnover during Orthodontic Tooth Movement. Am J Orthod Dentofacial Orthop 1993;103(4):320-6.	Exclusion by title
113	Kim SH, Choi YS, Hwang EH, Chung KR, Kook YA, Nelson G. Surgical positioning of orthodontic mini-implants with guides fabricated on models replicated with cone-beam computed tomography. Am J Orthod Dentofacial Orthop 2007;131(4):S82-S9.	Exclusion by title
114	Kisnisci RS, Iseri H, Tuz HH, Altug AT. Dentoalveolar distraction osteogenesis for rapid orthodontic canine retraction. Journal of Oral and Maxillofacial Surgery. 2002;60(4):389-94.	Exclusion by title
115	Kiukkonen A, Viluksela M, Sahlberg C, Alaluusua S, Tuomisto JT, Tuomisto J, et al. Response of the incisor tooth to 2,3,7,8-tetrachlorodibenzo-p-dioxin in a dioxin-resistant and a dioxin-sensitive rat strain. Toxicological Sciences. 2002;69(2):482-9.	Exclusion by title
116	Knosel M, Attin R, Becker K, Attin T. A randomized CIE L*a*b* evaluation of external bleaching therapy effects on fluorotic enamel stains. Quintessence Int. 2008;39(5):391-9.	Exclusion by title
117	Koenig-Maunsell R, Villanelo-Ninapaytan MS, Lavado-Torres A, Cieza-Quiroz CL, Herrera-Chavez CMdJ. La influencia del camino en el color, altura, ancho y altura del margen Gingival en una sonrisa percibida como atractiva. Kiru. 2013;10(2):116-32.	Exclusion by title
118	Krisdapong S, Prasertsom P, Rattananangsim K, Sheiham A. Associations between perceived needs for dental treatment, oral health-related quality of life and oral diseases in school-aged Thai children. Community Dentistry and Oral Epidemiology. 2014;42(4):323-32.	Exclusion by title
119	Lai HC, Zhang ZY, Wang F, Zhuang LF, Liu X, Pu YP. Evaluation of soft-tissue alteration around implant-supported single-tooth restoration in the anterior maxilla: the pink esthetic score. Clinical Oral Implants Research. 2008;19(6):560-4.	Exclusion by title
120	Lamb DJ, Ellis B, Patel R. Restorative treatment need assessment by dentally and nondentally trained subjects. J Oral Rehabil. 1998;25(1):34-9.	Exclusion by title
121	Lazzaretti DN, Bortoluzzi GS, Torres Fernandes LF, Rodriguez R, Grehs RA, Martins Hartmann MS. Histologic evaluation of human pulp tissue after orthodontic intrusion. J Endod. 2014;40(10):1537-40.	Exclusion by title
122	Learreta JA, Bono AE, Maffia G, Beas J. Identificación de patologías articulares por medio del estudio del ciclo masticatorio, en pacientes preortodóncicos. Parte I: descripción del ciclo masticatorio. Ortodoncia. 2003;67(133):8-13.	Exclusion by title
123	Lee RJ, Pham J, Choy M, Weissheimer A, Dougherty HL, Jr., Sameshima GT, et al. Monitoring of typodont root movement via crown superimposition of single cone-beam computed tomography and consecutive intraoral scans. Am J Orthod Dentofacial Orthop. 2014;145(3):399-409.	Exclusion by title
124	Lee RJ, Weissheimer A, Pham J, Go L, de Menezes LM, Redmond WR, et al. Three-dimensional monitoring of root movement during orthodontic treatment. Am J Orthod Dentofacial Orthop. 2015;147(1):132-42.	Exclusion by title
125	Lekic P, Kenny D, Moe HK, Barretti E, McCulloch CA. Relationship of clonogenic capacity to plating efficiency and vital dye staining of human periodontal ligament cells: implications for tooth replantation. J Periodontal Res. 1996;31(4):294-300.	Exclusion by title
126	Li N, Han B, Zhang Q. Antibacterial properties of inorganic nano-antibacterial materials in orthodontics. Chinese Journal of Tissue Engineering Research. 2015;19(12):1953-7.	Exclusion by title
127	Lim YK, Lee YK. Influence of light transmittance and background reflectance on the light curing of adhesives used to bond esthetic brackets. Am J Orthod Dentofacial Orthop 2007;132(1).	Exclusion by title
128	Ling KK, Ho CT, Kravchuk O, Olive RJ. Comparison of surgical and non-surgical methods of treating palatally impacted canines. II. Aesthetic outcomes. Aust Orthod J. 2007;23(1):8-15.	Exclusion by title
129	Liu X, Huang J, Bai YD, Wang XJ, Baker A, Chen FM, et al. Conservation of root-fractured primary teeth-report of a case. Dental Traumatology. 2013;29(6):498-501.	Exclusion by title
130	Loguercio AD, Amaral RCd, Garcia E, Reis A. Use of flowable composite as intermediary layer in non-cariou cervical lesions restored with composite resin: 48-month follow-up. Rev odonto ciênc. 2010;25(2):216-20.	Exclusion by title
131	Lundin SA, Rasmusson CG. Clinical evaluation of a resin composite and bonding agent in Class I and II restorations: 2-Year results. Quintessence International. 2004;35(9):758-62.	Exclusion by title
132	Mansilla J, Solis C, Chávez-Lomeli ME, Gama JE. Analysis of colored teeth from Precolumbian tlatelolco: Postmortem transformation or intravital processes? American Journal of Physical Anthropology. 2003;120(1):73-82.	Exclusion by title
133	Marchi JJ, Froner AM, Alves HL, Bergmann CP, Araujo FB. Analysis of primary tooth dentin after indirect pulp capping. J Dent Child (Chic). 2008;75(3):295-300.	Exclusion by title
134	Mariaca de Botero L, Botero Mariaca P. Diagnóstico y tratamiento temprano de la malposición intralveolar de caninos permanentes. Rev Fac Odontol Univ Antioq. 2002;13(2):21-9.	Exclusion by title
135	Marini I, Bortolotti F, Incerti Parenti S, Gatto MR, Alessandri Bonetti G. Combined effects of repeated oral hygiene motivation and type of toothbrush on orthodontic patients: A blind randomized clinical trial. Angle Orthodontist. 2014;84(5):896-901.	Exclusion by title
136	Martins C, Siqueira WL, Guimarães Primo LSS. Oral and salivary flow characteristics of a group of Brazilian children and adolescents with chronic renal failure. Pediatric Nephrology. 2008;23(4):619-24.	Exclusion by title
137	Martins CC, Feitosa NB, Vale MP, Paiva SM. Parents' perceptions of oral health conditions depicted in photographs of anterior permanent teeth. European Journal of Paediatric Dentistry. 2010;11(4):203-9.	Exclusion by title
138	Maurya R, Tripathi T, Rai P. New generation of color bonding: a comparative in vitro study. Indian J Dent Res. 2011;22(5):733-4.	Exclusion by title
139	Mejàre I, Bergman E, Grindefjord M. Hypomineralized molars and incisors of unknown origin: Treatment outcome at age 18 years. International Journal of Paediatric Dentistry. 2005;15(1):20-8.	Exclusion by title
140	Mejía PP, González A, Martínez Y, Villabón C, Sierra S, Gómez V. Manejo endodóntico conservador de un caso de Dens Invaginatus tipo III con vitalidad y lesión periapical. CES Odontol. 2009;22(1):31-8.	Exclusion by title

141	Mello CCd, Mazaro JVQ, Zavanelli AC, Zavanelli RA. Laminados cerâmicos para resolução estética de dentes manchados por tetraciclina. Full dent sci. 2012;4(13):186-94.	Exclusion by title
142	Meng CL, Li CH, Wang WN. Bond strength with APF applied after acid etching. Am J Orthod Dentofacial Orthop. 1998;114(5):510-3.	Exclusion by title
143	Millán Isea RE, Salinas Millán YJ, Maestre LP, Paz de Gudiño M. Enfermedad periodontal y tratamiento ortodóncico: reporte de un caso clínico. Acta Odontol Venez. 2007;45(2):280-2.	Exclusion by title
144	Modena KCdS, Colombini-ishikiriama BL, Ishikiriama SK, Scotti CK, Nahsan FPS. Multidisciplinary esthetic approach for multiple anterior diastema: 36-month clinical performance. Braz dent sci. 2015;18(4):110-9.	Exclusion by title
145	Montenegro-Arana A, Arana-Gordillo LA, Farana D, Davila-Sanchez A, Jadad E, Coelho U, et al. Randomized Double-blind Clinical Trial of Bleaching Products in Patients Wearing Orthodontic Devices. Operative Dentistry. 2016;41(4):379-87.	Exclusion by title
146	Moreira KMS, Rezende CP, Silva DV, Imperato JCP, Navarro RS, Reis JB. Reabilitação do incisivo central decíduo fusionado ao supranumerário: relato de caso clínico. Rev Assoc Paul Cir Dent. 2016;70(3):302-5.	Exclusion by title
147	Moura-Netto C, Guglielmi CdAB, Mello-Moura ACV, Caldeira CL, Raggio DP, Palo RM. Nd:YAG laser irradiation effect on apical intracanal dentin - a microleakage and SEM evaluation. Braz Dent J. 2011;22(5):377-81.	Exclusion by title
148	Mussa R, Esposito SJ, Cowper TR. The use of colored elastomeric "O"s as a motivational instrument for patients with anodontia: Report of case. Journal of Dentistry for Children. 1999;66(2):98-+.	Exclusion by title
149	Müssig E, Lux CJ, Staehle HJ, Stellzig-Eisenhauer A, Komposch G. Applications for direct composite restorations in orthodontics. J Orofac Orthop 2004;65(2):164-79.	Exclusion by title
150	Nakajima K. A study of primary bone grafting for unilateral complete cleft lip and palate (Japanese). Acta Medica (Igaku Kenkyu). 1974;44(1):17-41.	Exclusion by title
151	Nakasima A, Hu JR, Ichinose M, Shimada H. Potential Application of Shape Memory Plastic as Elastic-Material in Clinical Orthodontics. Eur J Orthod 1991;13(3):179-86.	Exclusion by title
152	Namura Y, Tsuruoka T, Ryu C, Kaketani M, Shimizu N. Usefulness of orthodontic adhesive-containing fluorescent dye. Eur J Orthod 2010;32(6):620-6.	Exclusion by title
153	Naorungroj S. Esthetic Reconstruction of Diastema with Adhesive Tooth-Colored Restorations and Hyaluronic Acid Fillers. Case Reports in Dentistry. 2017.	Exclusion by title
154	Nelson PA, Artun J. Alveolar bone loss of maxillary anterior teeth in adult orthodontic patients. Am J Orthod Dentofacial Orthop 1997;111(3):328-34.	Exclusion by title
155	Neumann P, Bourauel C, Jäger A. Corrosion and permanent fracture resistance of coated and conventional orthodontic wires. Journal of Materials Science: Materials in Medicine. 2002;13(2):141-7.	Exclusion by title
156	Nie EM, Jiang R, Zhang CY, Zeng JD, Tan JZ. Temporary protection of cracked teeth using orthodontic band and resin temporary crown. Chinese Journal of Tissue Engineering Research. 2015;19(30):4805-9.	Exclusion by title
157	Nocca G, Chimenti C, Parziale V, Gambarini G, Giardina B, Lupi A. In vitro comparison of the cytotoxicity of two orthodontic composite resins. Minerva Stomatol. 2006;55(5):297-305.	Exclusion by title
158	Oesterle LJ, Shellhart WC, Belanger GK. Effect of tacking time on bond strength of light-cured adhesives. J Clin Orthod. 1997;31(7):449-53.	Exclusion by title
159	Oesterle LJ, Shellhart WC, Belanger GK. The use of bovine enamel in bonding studies. Am J Orthod Dentofacial Orthop. 1998;114(5):514-9.	Exclusion by title
160	Olmedo DG, Nalli G, Verdu S, Paparella ML, Cabrini RL. Exfoliative cytology and titanium dental implants: a pilot study. J Periodontol. 2013;84(1):78-83.	Exclusion by title
161	Ovrebo RC, Raadal M. Microleakage in fissures sealed with resin or glass ionomer cement. Scand J Dent Res. 1990;98(1):66-9.	Exclusion by title
162	Patel S, Fanshawe T, Bister D, Cobourne MT. Survival and success of maxillary canine autotransplantation: a retrospective investigation. Eur J Orthod 2011;33(3):298-304.	Exclusion by title
163	Pedrazzi H, Andrade MFd, Takeuchi CYG, Santos CRd. Reabilitação estética do sorriso por meio do sistema cerâmico dissilicato de lítio: relato de caso. Full dent sci. 2014;6(21):112-7.	Exclusion by title
164	Peña Ruiz T, Martínez Brito I, Delgado Ramos A. Tratamiento integrado ortoperiodontal en la periodontitis juvenil. Presentación de un caso. Rev Cubana Estomatol. 2007;44(4).	Exclusion by title
165	Peng Y, Wu R, Qu W, Wu W, Chen J, Fang J, et al. Effect of visual method vs plaque disclosure in enhancing oral hygiene in adolescents and young adults: a single-blind randomized controlled trial. Am J Orthod Dentofacial Orthop. 2014;145(3):280-6.	Exclusion by title
166	Perea Paz M, Aliaga Del Castillo A, Aliaga Del Castillo R. Exodoncia de diente supernumerario, canino en transposición y enucleación de odontoma complejo: reporte de caso. Rev estomatol Hered. 2010;20(2):90-3.	Exclusion by title
167	Pereira BR, Tanaka OM, Lima AA, Guariza-Filho O, Maruo H, Camargo ES. Metal and ceramic bracket effects on human buccal mucosa epithelial cells. Angle Orthod. 2009;79(2):373-9.	Exclusion by title
168	Peres KG, Frazão P, Roncalli AG. Padrão epidemiológico das oclusopatias muito graves em adolescentes brasileiros. Rev Saude Publica. 2013;47(supl.3):109-17.	Exclusion by title
169	Pérez Araya J, González JP, González Valdenegro R, Lorenzo Vladilo M. Necrotizing cervical fasciitis, a review of the literature in association to a clinical case. International Journal of Oral and Maxillofacial Surgery. 2011;40(10):1184.	Exclusion by title
170	Pérez Araya J, González JP, González Valdenegro R. Oral and radiological characteristics of gardner syndrome patients: A report on three cases of members of the same family. International Journal of Oral and Maxillofacial Surgery. 2011;40(10):1142.	Exclusion by title
171	Picinato-Pirola MN, Mestriner Junior W, Mello-Filho F, Trawitzki L. Masticatory efficiency in class III dentofacial deformities before and after orthognathic surgery. Journal of Oral and Maxillofacial Surgery. 2011;69(9):e81-e2.	Exclusion by title
172	Poosti M, Ahrari F, Moosavi H, Najjaran H. The effect of fractional CO2 laser irradiation on remineralization of enamel white spot lesions. Lasers in Medical Science. 2014;29(4):1349-55.	Exclusion by title
173	Post LK, Lima FG, Gerhardt-Oliveira M, Demarco FF, Xavier CB. Sealing ability of MTA and amalgam in different root-end preparations and resection bevel angles: an in vitro evaluation using marginal dye leakage. Braz Dent J. 2010;21(5):416-9.	Exclusion by title
174	Premaraj T, Simet S, Beatty M, Premaraj S. Oral epithelial cell reaction after exposure to Invisalign plastic material. Am J Orthod Dentofacial Orthop. 2014;145(1):64-71.	Exclusion by title
175	Rai P, Pandey RK, Khanna R. A multidisciplinary approach to the management of traumatic intrusion in immature permanent teeth. BMJ Case Reports. 2016;2016.	Exclusion by title
176	Robertson A, Andreasen FM, Bergenholtz G, Andreasen JO, Noren JG. Incidence of pulp necrosis subsequent to pulp canal obliteration from trauma of permanent incisors. Journal of Endodontics. 1996;22(10):557-60.	Exclusion by title
177	Robertsson S, Mohlin B, Thilander B. Aesthetic evaluation in subjects treated due to congenitally missing maxillary laterals. A comparison of perception in patients, parents and dentists. Swed Dent J. 2010;34(4):177-86.	Exclusion by title

178	Rohilla M, Pandit IK, Srivastava N. Why the carioprotective potential of luting cements crucial? J Indian Soc Pedod Prev Dent. 2009;27(4):219-23.	Exclusion by title
179	Rongo R, Valletta R, Buccia R, Riviuccio V, Galeotti A, Michelotti A, et al. In vitro biocompatibility of nickel-titanium esthetic orthodontic archwires. Angle Orthodontist. 2016;86(5):789-95.	Exclusion by title
180	Salehi P, Momeni Danaie S. Comparison of the antibacterial effects of persica mouthwash with chlorhexidine on streptococcus mutans in orthodontic patients. Daru. 2006;14(4):178-82.	Exclusion by title
181	Salmassian R, Oesterle LJ, Shellhart WC, Newman SM. Comparison of the efficacy of ibuprofen and acetaminophen in controlling pain after orthodontic tooth movement. Am J Orthod Dentofacial Orthop. 2009;135(4):516-21.	Exclusion by title
182	Samorodnitsky-Naveh GR, Geiger SB, Levin L. Patients' satisfaction with dental esthetics. Journal of the American Dental Association. 2007;138(6):805-8.	Exclusion by title
183	Sandhu M, Gulia S, Nagpal M, Sachdev V. Circular enamel hypoplasia: A rare enamel developmental disturbance in permanent teeth. Journal of Clinical and Diagnostic Research. 2014;8(8):ZD39-ZD40.	Exclusion by title
184	Santiago SL, Passos VF, Navarro MFdL, Lauris JRP, Vieira AHM, Franco EB. Two-year clinical evaluation of resinous restorative systems in non-carious cervical lesions. Braz Dent J. 2010;21(3):229-34.	Exclusion by title
185	Santos BM, Pithon MM, Ruellas ACD, Sant'Anna EF. Shear bond strength of brackets bonded with hydrophilic and hydrophobic bond systems under contamination. Angle Orthodontist. 2010;80(5):963-7.	Exclusion by title
186	Sarul M, Kawala B, Kawala M, Antoszewska-Smith J. Do the NiTi low and constant force levels remain stable in vivo? Eur J Orthod. 2015;37(6):656-64.	Exclusion by title
187	Sawicki L, Pameijer CH, Emerich K, Adamowicz-Klepalska B. Histological evaluation of mineral trioxide aggregate and calcium hydroxide in direct pulp capping of human immature permanent teeth. Am J Dent. 2008;21(4):262-6.	Exclusion by title
188	Schatzle M, Imfeld T, Sener B, Schmidlin PR. In vitro tooth cleaning efficacy of manual toothbrushes around brackets. Eur J Orthod. 2009;31(1):103-7.	Exclusion by title
189	Schatzle M, Sener B, Schmidlin PR, Imfeld T, Attin T. In vitro tooth cleaning efficacy of electric toothbrushes around brackets. Eur J Orthod. 2010;32(5):481-9.	Exclusion by title
190	Schmidlin PR, Schug J. [Profilometric determination of substance removal and surface morphology after microabrasion in vivo]. Schweiz Monatsschr Zahnmed. 2001;111(12):1425-34.	Exclusion by title
191	Schmidt CJ, Tatum SA. Cosmetic dentistry. Current Opinion in Otolaryngology and Head and Neck Surgery. 2006;14(4):254-9.	Exclusion by title
192	Schneider O. Cerec Veneers - Practical Procedure and Case Presentation. International Journal of Computerized Dentistry. 2003;6(3):283-92.	Exclusion by title
193	Shafiq MK, Mirza AJ, Ahmed SZ. Management of crown-root fracture of an incisor using a prefabricated metal post. Pakistan Journal of Medical Sciences. 2011;27(2):434-7.	Exclusion by title
194	Shahabi M, Jahanbin A, Esmaily H, Sharifi H, Salari S. Comparison of Some Dietary Habits on Corrosion Behavior of Stainless Steel Brackets: An in vitro Study. Journal of Clinical Pediatric Dentistry. 2011;35(4):429-32.	Exclusion by title
195	Shan LH, Cui ZQ, Shen QH, Gao Q, Qiu ZX. Application of light-cure resin-modified glass ionomer cement in orthodontic practice. Journal of Clinical Rehabilitative Tissue Engineering Research. 2008;12(6):1149-52.	Exclusion by title
196	Sharma R, Kumar P, Bhargava N, Sharma AK, Srivastava S, Jain S, et al. Dental and skeletal fluorosis: A review. Medico-Legal Update. 2013;13(2):151-5.	Exclusion by title
197	Shyagali TR, Bhayya DP, Subramaniam S, Urs CB. Finite element study on modification of bracket base and its effects on bond strength. Dental Press J Orthod. 2015;20(2):76-82.	Exclusion by title
198	Silva GA, Gava E, Lanza LD, Estrela C, Alves JB. Subclinical failures of direct pulp capping of human teeth by using a dentin bonding system. J Endod. 2013;39(2):182-9.	Exclusion by title
199	Silva W, Olímpio L, Frederico P, Duarte PS, Pinto T, Montenegro G. Reabilitação oral total com materiais adesivos: construindo um protocolo previsível. Full dent sci. 2012;4(13):100-7.	Exclusion by title
200	Singh HP, Shetty S, Patil P, Sethi N, Singh A, Raghunandan BN. Correlation between collagen fibers and radiographic patterns of keratocystic odontogenic tumour. Journal of Clinical and Diagnostic Research. 2014;8(8):ZC16-ZC8.	Exclusion by title
201	Singh K, Kumar N, Choudhary N, Gupta N. Unconventional prosthodontics for the aesthetic rehabilitation of discoloured rotated maxillary central incisor. BMJ Case Reports. 2013.	Exclusion by title
202	Smailiene D, Kavaliauskiene A, Pacauskiene I. Posttreatment status of palatally impacted maxillary canines treated applying 2 different surgical-orthodontic methods. Medicina (Kaunas). 2013;49(8):354-60.	Exclusion by title
203	Smiech-Slomkowska G, Jablonska-Zrobek J. The effect of oral health education on dental plaque development and the level of caries-related Streptococcus mutans and Lactobacillus spp. Eur J Orthod. 2007;29(2):157-60.	Exclusion by title
204	Solem RC, Marasco R, Guitierrez-Pulido L, Nielsen I, Kim SH, Nelson G. Three-dimensional soft-tissue and hard-tissue changes in the treatment of bimaxillary protrusion. Am J Orthod Dentofacial Orthop. 2013;144(2):218-28.	Exclusion by title
205	Stritikus J, Owens B. An in vitro study of microleakage of occlusal composite restorations polymerized by a conventional curing light and a PAC curing light. J Clin Pediatr Dent. 2000;24(3):221-7.	Exclusion by title
206	Subay RK, Kayatas M, Canikloglu C. Delayed multidisciplinary management of an extrusively luxated maxillary central incisor. Dental Traumatology. 2007;23(2):82-4.	Exclusion by title
207	Succaria F, Morgano SM. Prescribing a dental ceramic material: Zirconia vs lithium-disilicate. Saudi Dental Journal. 2011;23(4):165-6.	Exclusion by title
208	Surgeon TW. Fundamentals of small animal orthodontics. Veterinary Clinics of North America - Small Animal Practice. 2005;35(4 SPEC. ISS.):869-89.	Exclusion by title
209	Teixeira L, Pereira Bdo R, Bortoly TG, Brancher JA, Tanaka OM, Guariza-Filho O. The environmental influence of Light Coke, phosphoric acid, and citric acid on elastomeric chains. J Contemp Dent Pract. 2008;9(7):17-24.	Exclusion by title
210	Thelen DS, Trovik TA, Bardsen A. Impact of traumatic dental injuries with unmet treatment need on daily life among Albanian adolescents: a case-control study. Dent Traumatol. 2011;27(2):88-94.	Exclusion by title
211	Thomas BW, Hook CR, Draughn RA. Laser-aided degradation of composite resin. Angle Orthod. 1996;66(4):281-6.	Exclusion by title
212	Thordarson A, Zachrisson BU, Mjor IA. Remodeling of Canines to the Shape of Lateral Incisors by Grinding - a Long-Term Clinical and Radiographic Evaluation. Am J Orthod Dentofacial Orthop. 1991;100(2):123-32.	Exclusion by title
213	Tian CS, He P, Song JL, Wu P, Deng F, Gao X. The typodont model based investigation on three-dimensional movement of incisors in alignment stage when applying MDD and MBT orthodontic brackets. Chinese Journal of Biomedical Engineering. 2012;31(4):494-501.	Exclusion by title
214	Toksavul S, Turkun M, Toman M. Esthetic enhancement of ceramic crowns with zirconia dowels and cores: A clinical report. Journal of Prosthetic Dentistry. 2004;92(2):116-9.	Exclusion by title

215	Torchi SdO, Freitas KMSd, Lima NCJ, Valarelli FP, Cançado RH. Comparação das alterações esqueléticas, dentárias e tegumentares entre pacientes leucodermas e melanodermas tratados ortodonticamente com extrações. Ortho Sci, Orthod sci pract. 2016;9(33):39-47.	Exclusion by title
216	Torlakovic L, Olsen I, Petzold C, Tiainen H, Ogaard B. Clinical color intensity of white spot lesions might be a better predictor of enamel demineralization depth than traditional clinical grading. Am J Orthod Dentofacial Orthop 2012;142(2):191-8.	Exclusion by title
217	Tosheska-Trajkovska K, Biljali S, Nikolov G, Spasov Z, Toseska-Spasova N. Eating disorders in orthodontic practice and associated laboratory findings. Clinical Chemistry and Laboratory Medicine. 2014;52:S1296.	Exclusion by title
218	Turkkahraman H, Adanir N, Gungor AY, Alkis H. In vitro evaluation of shear bond strengths of colour change adhesives. Eur J Orthod. 2010;32(5):571-4.	Exclusion by title
219	Twetman S, McWilliam JS, Hallgren A, Oliveby A. Cariostatic effect of glass ionomer retained orthodontic appliances. An in vivo study. Swed Dent J. 1997;21(5):169-75.	Exclusion by title
220	Ueda H, Almeida FR, Lowe AA, Ruse ND. Changes in occlusal contact area during oral appliance therapy assessed on study models. Angle Orthod. 2008;78(5):866-72.	Exclusion by title
221	Uysal T, Ramoglu SI, Ertas H, Ulker M. Microleakage of orthodontic band cement at the cement-enamel and cement-band interfaces. Am J Orthod Dentofacial Orthop. 2010;137(4):534-9.	Exclusion by title
222	Villarreal E, Sánchez Soler LA, Masip S, Espías Gómez Á. Manejo estratificado de composites en sector anterior: un gran paso en odontología mínimamente invasiva. Dentum (Barc). 2007;7(3):118-23.	Exclusion by title
223	Wakasa K, Ikeda A, Yoshida Y, Yamaki M. Dental castable glass ceramics: Ceramming treatment and colour property. Journal of Materials Science: Materials in Medicine. 1995;6(1):29-31.	Exclusion by title
224	Wang DF, Jia S, Ye RR, Bao XJ. Restoration of clearance between maxillary central incisors by everStick® fibre reinforced composite: a case report. Journal of Shanghai Jiaotong University (Medical Science). 2015;35(9):1426-30.	Exclusion by title
225	Winik R, Araki ÂT, Negrão JAA, Bello-Silva MS, Lage-Marques JL. Sealer penetration and marginal permeability after apicoectomy varying retrocavity preparation and retrofilling material. Braz Dent J. 2006;17(4):323-7.	Exclusion by title
226	Wriedt S, Martin M, Al-Nawas B, Wehrbein H. Long-term Effects of Traumatic Injuries to Incisors and Periodontal Tissues during Childhood A Retrospective Study. Journal of Orofacial Orthopedics-Fortschritte Der Kieferorthopadie. 2010;71(5):318-29.	Exclusion by title
227	Wudel J, Valdez S, Perkins JN, Liang C, Blanchard A, Allen GC. Parent, dentist, and orthodontist satisfaction following alveolar cleft repair using recombinant human bone morphogenic protein. Cleft Palate-Craniofacial Journal. 2015;52(3):e41-e6.	Exclusion by title
228	Xie GL, Meng YK. A Case Report of Esthetic and Functional Correction of Maxillary Protrusion Using a Prosthodontics-Centered Multidisciplinary Approach. International Journal of Periodontics & Restorative Dentistry. 2016;36(2):E33-E40.	Exclusion by title
229	Yi I, Chan KH, Tsuji GH, Staninec M, Darling CL, Fried D. Selective removal of esthetic composite restorations with spectral guided laser ablation. Lasers in Dentistry Xxii. Proceedings of SPIE. 96922016.	Exclusion by title
230	Yoshikawa M, Kakei M, Fujita M, Tanaka A, Suda N. Misconceptions regarding tooth whitening by hydrogen peroxide. Bone. 2011;48:S116.	Exclusion by title
231	Zarrabi MH, Javidi M, Jafarian AH, Joushan B. Immunohistochemical expression of fibronectin and tenascin in human tooth pulp capped with mineral trioxide aggregate and a novel endodontic cement. J Endod. 2011;37(12):1613-8.	Exclusion by title
232	Zulian F, Cuffaro G, Sperotto F. Scleroderma in children: An update. Current Opinion in Rheumatology. 2013;25(5):643-50.	Exclusion by title
233	Zyla T, Kawala B, Antoszewska-Smith J, Kawala M. Black stain and dental caries: A review of the literature. BioMed Research International. 2015;2015.	Exclusion by title
234	Armstrong D, Shen G, Petocz P, Darendeliler MA. Excess adhesive flash upon bracket placement. A typodont study comparing APC PLUS and Transbond XT. Angle Orthod. 2007;77(6):1101-8.	Exclusion by abstract
235	Barnes CM, Russell CM, Gerbo LR, Wells BR, Barnes DW. Effects of an air-powder polishing system on orthodontically bracketed and banded teeth. Am J Orthod Dentofacial Orthop. 1990;97(1):74-81.	Exclusion by abstract
236	Benham AW, Campbell PM, Buschang PH. Effectiveness of pit and fissure sealants in reducing white spot lesions during orthodontic treatment. A pilot study. Angle Orthod. 2009;79(2):338-45.	Exclusion by abstract
237	Chapman JL. Bond failure rates of two self-ligating brackets: a randomised clinical trial. Aust Orthod J. 2011;27(2):139-44.	Exclusion by abstract
238	Chetpakdeehtit W, Wahss J, Woo T, Hugander M, Mohlin B, Hagberg C. Esthetic views on facial and dental appearance in young adults with treated bilateral cleft lip and palate (BCLP). A comparison between professional and non-professional evaluators. Swed Dent J. 2011;35(3):151-7.	Exclusion by abstract
239	Doshi UH, Bhad-Patil WA. Static frictional force and surface roughness of various bracket and wire combinations. Am J Orthod Dentofacial Orthop. 2011;139(1):74-9.	Exclusion by abstract
240	Elayyan F, Silikas N, Bearn D. Ex vivo surface and mechanical properties of coated orthodontic archwires. Eur J Orthod. 2008;30(6):661-7.	Exclusion by abstract
241	Hu B, Song J, Wu P, He P, Deng F, Gao X. [A comparative study on three-dimensional movement of anterior teeth between applying MDD appliances and applying three common fixed appliances in the initial alignment stage]. Sheng Wu Yi Xue Gong Cheng Xue Za Zhi. 2013;30(5):1039-43.	Exclusion by abstract
242	Kim S, Kim EY, Jeong TS, Kim JW. The evaluation of resin infiltration for masking labial enamel white spot lesions. Int J Paediatr Dent. 2011;21(4):241-8.	Exclusion by abstract
243	Knosel M, Eckstein A, Helms HJ. Durability of esthetic improvement following Icon resin infiltration of multibracket-induced white spot lesions compared with no therapy over 6 months: a single-center, split-mouth, randomized clinical trial. Am J Orthod Dentofacial Orthop. 2013;144(1):86-96.	Exclusion by abstract
244	Moosavi H, Ahrari F, Mohamadipour H. The effect of different surface treatments of demineralised enamel on microleakage under metal orthodontic brackets. Prog Orthod. 2013;14:2.	Exclusion by abstract
245	Nagpal R, Tewari S, Gupta R. Effect of various surface treatments on the microleakage and ultrastructure of resin-tooth interface. Oper Dent. 2007;32(1):16-23.	Exclusion by abstract
246	Restrepo M, Bussaneli DG, Jeremias F, Cordeiro RC, Raveli DB, Magalhaes AC, et al. Control of White Spot Lesions with Use of Fluoride Varnish or Chlorhexidine Gel During Orthodontic Treatment A Randomized Clinical Trial. J Clin Pediatr Dent. 2016;40(4):274-80.	Exclusion by abstract
247	Rudge P, Sherriff M, Bister D. A comparison of roughness parameters and friction coefficients of aesthetic archwires. Eur J Orthod. 2015;37(1):49-55.	Exclusion by abstract
248	Uysal T, Ulker M, Ramoglu SI, Ertas H. Microleakage under metallic and ceramic brackets bonded with orthodontic self-etching primer systems. Angle Orthod. 2008;78(6):1089-94.	Exclusion by abstract
249	Voudouris JC. Interactive edgewise mechanisms: form and function comparison with conventional edgewise brackets. Am J Orthod Dentofacial Orthop. 1997;111(2):119-40.	Exclusion by abstract
250	Yagci A, Uysal T, Ulker M, Ramoglu SI. Microleakage under orthodontic brackets bonded with the custom base indirect bonding technique. Eur J Orthod. 2010;32(3):259-63.	Exclusion by abstract

251	Bezerra GL, Torres CRG, Tonetto MR, Borges AH, Kuga MC, Bandeca MC, et al. Shear Bond Strength of Orthodontic Brackets Fixed with Remineralizing Adhesive Systems after Simulating One Year of Orthodontic Treatment. <i>Scientific World Journal</i> . 2015;2015.	Exclusion by abstract
252	Elnafar AAS, Alam MK, Hassan R, Purnal K. Enamel surface preparations and shear bond strength of orthodontic brackets: A review. <i>International Medical Journal</i> . 2015;22(3):194-8.	Exclusion by abstract
253	Georgieva K, Katreva I, Abadjiev M. Mastication, phonetics and esthetics as a final result of partial or complete denture treatment. <i>Journal of IMAB - Annual Proceeding (Scientific Papers)</i> . 2015;21(3):846-8.	Exclusion by abstract
254	Goldfogel M, Noonan Jr JE. A non-surgical approach to an anterior open bite: Report of a case. <i>Gerodontology</i> . 1986;2(1):39-42.	Exclusion by abstract
255	Rabchinsky DS, Powers JM. Color stability and stain resistance of direct-bonding orthodontic cements. <i>American Journal of Orthodontics</i> . 1979;76(2):170-7.	Exclusion by abstract
256	Wriedt S, Schepke U, Wehrbein H. The discoloring effects of food on the color stability of esthetic brackets - An in-vitro study. <i>J Orofasc Orthop</i> 2007;68(4):308-20.	Exclusion by abstract
257	Knösel M, Reus M, Rosenberger A, Ziebolz D. A novel method for testing the veridicality of dental colour assessments. <i>Eur J Orthod</i> 2012;34(1):19-24.	Exclusion by abstract
258	Alkhatib MN, Holt R, Bedi R. Prevalence of self-assessed tooth discolouration in the United Kingdom. <i>J Dent</i> 2004;32(7):561-6.	Exclusion by abstract
259	Hallgren K, Akyalcin S, English J, Tufekci E, Paravina RD. Color Properties of Demineralized Enamel Surfaces Treated with a Resin Infiltration System. <i>Journal of Esthetic and Restorative Dentistry</i> . 2016;28(5):339-46.	Exclusion by abstract
260	Lee YK. Changes in the reflected and transmitted color of esthetic brackets after thermal cycling. <i>Am J Orthod Dentofacial Orthop</i> 2008;133(5).	Exclusion by abstract
261	Lee YK. Colour and translucency of tooth-coloured orthodontic brackets. <i>Eur J Orthod</i> 2008;30(2):205-10.	Exclusion by abstract
262	Leland A, Akyalcin S, English JD, Tufekci E, Paravina R. Evaluation of staining and color changes of a resin infiltration system. <i>Angle Orthodontist</i> . 2016;86(6):900-4.	Exclusion by abstract
263	Lopes H, Maia LH, Araujo MV, Elias CN, Ruellas ACO. Colour stability of aesthetic brackets: ceramic and plastic. <i>Australian Orthodontic Journal</i> . 2013;29(1):13-20.	Exclusion by abstract
264	Qin YJ, Zhang GD, Zhang Y, Ping YF, Zhao CY. Natural reversal of tooth discoloration and pulpal response to testing following removal of a miniscrew implant for orthodontic anchorage: a case report. <i>International Endodontic Journal</i> . 2016;49(4):402-9.	Exclusion by abstract
265	Cano Lavín V, Díaz Christians L, López Martínez R. Daños a tejidos dentarios y peridentarios utilizando la técnica ortodóncica de Poussin. <i>Rev Cubana Estomatol</i> . 1994;31(1):6-10.	Exclusion by abstract
266	García EJ, Serrano APM. Tratamiento estético de manchas blancas de esmalte posortodoncia: asociación de blanqueamiento dental e infiltración resinosa. <i>Rev Asoc Odontol Argent</i> . 2013;101(1):17-20.	Exclusion by abstract
267	Romano FL, Valério RA, Gomes-Silva JM, Ferreira JTL, Faria G, Borsatto MC. Clinical evaluation of the failure rate of metallic brackets bonded with orthodontic composites. <i>Braz Dent J</i> . 2012;23(4):399-402.	Exclusion by abstract
268	Eckstein A, Helms HJ, Knosel M. Camouflage effects following resin infiltration of postorthodontic white-spot lesions in vivo: One-year follow-up. <i>Angle Orthod</i> . 2015;85(3):374-80.	Excluded; not relevant
269	Gwinnett AJ. Corrosion of resin-bonded orthodontic brackets. <i>Am J Orthod</i> . 1982;81(6):441-6.	Excluded; not relevant
270	Hammad SM, El Banna M, El Zayat I, Mohsen MA. Effect of resin infiltration on white spot lesions after debonding orthodontic brackets. <i>Am J Dent</i> . 2012;25(1):3-8.	Excluded; not relevant
271	Jadad E, Montoya J, Arana G, Gordillo LA, Palo RM, Loguercio AD. Spectrophotometric evaluation of color alterations with a new dental bleaching product in patients wearing orthodontic appliances. <i>Am J Orthod Dentofacial Orthop</i> . 2011;140(1):e43-7.	Excluded; not relevant
272	Kappert HF, Jonas I, Rakosi T. [Importance of the corrosion factor in bracket adhesive technics]. <i>Fortschr Kieferorthop</i> . 1984;45(4):271-83.	Excluded; not relevant
273	Knosel M, Attin R, Becker K, Attin T. External bleaching effect on the color and luminosity of inactive white-spot lesions after fixed orthodontic appliances. <i>Angle Orthod</i> . 2007;77(4):646-52.	Excluded; not relevant
274	Kulas A, Illge C, Bekes K, Eckert AW, Fuhrmann RA, Hirsch C. Structural color changes in permanent enamel of patients with cleft lip and palate: a case-control study. <i>J Orofasc Orthop</i> . 2016;77(1):45-51.	Excluded; not relevant
275	Lopes Filho H, Maia LE, Araujo MV, Ruellas AC. Influence of optical properties of esthetic brackets (color, translucence, and fluorescence) on visual perception. <i>Am J Orthod Dentofacial Orthop</i> . 2012;141(4):460-7.	Excluded; not relevant
276	Millett DT, McCluskey LA, McAuley F, Creanor SL, Newell J, Love J. A comparative clinical trial of a compomer and a resin adhesive for orthodontic bonding. <i>Angle Orthod</i> . 2000;70(3):233-40.	Excluded; not relevant
277	Ajami S, Pakshir H, Babanouri N. Impact of nanohydroxyapatite on enamel surface roughness and color change after orthodontic debonding. <i>Prog Orthod</i> 2017;17(1).	Excluded; not relevant
278	Brascher AK, Zuran D, Feldmann RE, Benrath J. Patient survey on Invisalign(A (R)) treatment comparen the SmartTrack(A (R)) material to the previous aligner material. <i>Journal of Orofacial Orthopedics-Fortschritte Der Kieferorthopadie</i> . 2016;77(6):432-8.	Excluded; not relevant
279	Karamouzou A, Athanasiou AE, Papadopoulos MA. Tooth-color assessment after orthodontic treatment: The fundamental role of evidence-based medicine response. <i>Am J Orthod Dentofacial Orthop</i> 2011;139(3):291-.	Excluded; not relevant
280	Mangano A, Mangano A. Tooth-color assessment after orthodontic treatment: The fundamental role of evidence-based medicine. <i>Am J Orthod Dentofacial Orthop</i> 2011;139(3):290-1.	Excluded; not relevant
281	Sundfeld RH, Rahal V, Croll TP, De Aalexandre RS, Briso ALF. Enamel microabrasion followed by dental bleaching for patients after orthodontic treatment - Case reports. <i>Journal of Esthetic and Restorative Dentistry</i> . 2007;19(2):71-7.	Excluded; not relevant
282	Boncuk Y, Cehreli ZC, Polat-Ozsoy O. Effects of different orthodontic adhesives and resin removal techniques on enamel color alteration. <i>Angle Orthod</i> . 2014;84(4):634-41.	Excluded; not clinical study
283	Eliades T, Kakaboura A, Eliades G, Bradley TG. Comparison of enamel colour changes associated with orthodontic bonding using two different adhesives. <i>Eur J Orthod</i> . 2001;23(1):85-90.	Excluded; not clinical study
284	Trakyalı G, Ozdemir FI, Arun T. Enamel colour changes at debonding and after finishing procedures using five different adhesives. <i>Eur J Orthod</i> . 2009;31(4):397-401.	Excluded; not clinical study
285	Zaher AR, Abdalla EM, Abdel Motie MA, Rehman NA, Kassem H, Athanasiou AE. Enamel colour changes after debonding using various bonding systems. <i>J Orthod</i> . 2012;39(2):82-8.	Excluded; not clinical study
286	Jahanbin A, Ameri H, Khaleghimoghaddam R. Effect of adhesive types on enamel discolouration around orthodontic brackets. <i>Australian Orthodontic Journal</i> . 2009;25(1):19-23.	Excluded; not clinical study
287	Joo HJ, Lee YK, Lee DY, Kim YJ, Lim YK. Influence of orthodontic adhesives and clean-up procedures on the stain susceptibility of enamel after debonding. <i>Angle Orthodontist</i> . 2011;81(2):334-40.	Excluded; not clinical study
288	Wriedt S, Keller S, Wehrbein H. The effect of debonding and/or bleaching on enamel color - an in-vitro study. <i>Journal of Orofacial Orthopedics-Fortschritte Der Kieferorthopadie</i> . 2008;69(3):169-76.	Excluded; not clinical study
289	Chen QS, Zheng XC, Chen WT, Ni ZY, Zhou Y. Influence of orthodontic treatment with fixed appliances on enamel color: a systematic review. <i>Bmc Oral Health</i> . 2015;15.	Excluded; systematic review

290	Oltramari-Navarro PV, Titarelli JM, Marsicano JA, Henriques JF, Janson G, Lauris JR, et al. Effectiveness of 0.50% and 0.75% chlorhexidine dentifrices in orthodontic patients: a double-blind and randomized controlled trial. <i>Am J Orthod Dentofacial Orthop</i> . 2009;136(5):651-6.	Excluded; tooth-discoloring adjuncts
291	Olympio KP, Bardal PA, de MBJR, Buzalaf MA. Effectiveness of a chlorhexidine dentifrice in orthodontic patients: a randomized-controlled trial. <i>J Clin Periodontol</i> . 2006;33(6):421-6.	Excluded; tooth-discoloring adjuncts
292	Lee UL, Lee EJ, Seo HY, Han SH, Choi WC, Choi YJ. Prevalence and risk factors of tooth discoloration after orthognathic surgery: a retrospective study of 1455 patients. <i>International Journal of Oral and Maxillofacial Surgery</i> . 2016;45(11):1464-70.	Excluded; orthognathic treatment
293	Choi NC, Park YC, Jo YM, Lee KJ. Combined use of miniscrews and clear appliances for the treatment of bialveolar protrusion without conventional brackets. <i>Am J Orthod Dentofacial Orthop</i> 2009;135(5):671-81.	Excluded; color not assessed
294	Prihastuti Jufri Y. Relationship between fixed orthodontic treatment and Tooth Discoloration. Dissertation, Hasanuddin University, 2017.	Excluded; cross-sectional study
295	Anderson GB, Bowden J, Morrison EC, Caffesse RG. Clinical effects of chlorhexidine mouthwashes on patients undergoing orthodontic treatment. <i>Am J Orthod Dentofacial Orthop</i> . 1997;111(6):606-12.	Excluded; color assessed only mid-treatment
296	Al Maaitah EF, Abu Omar AA, Al-Khateeb SN. Effect of fixed orthodontic appliances bonded with different etching techniques on tooth color: a prospective clinical study. <i>Am J Orthod Dentofacial Orthop</i> . 2013;144(1):43-9.	Included
297	Corekci B, Toy E, Ozturk F, Malkoc S, Ozturk B. Effects of contemporary orthodontic composites on tooth color following short-term fixed orthodontic treatment: a controlled clinical study. <i>Turk J Med Sci</i> . 2015;45(6):1421-8.	Included
298	Karamouzos A, Athanasiou AE, Papadopoulos MA, Kolokithas G. Tooth-color assessment after orthodontic treatment: a prospective clinical trial. <i>Am J Orthod Dentofacial Orthop</i> . 2010;138(5):537.e1-8; discussion -9.	Included
299	Al-Laban YRA. Comparison of Enamel Color Alteration between Bonded and Free Unbonded Surfaces of Maxillary Anterior Teeth after Fixed Orthodontic Therapy : A Prospective Clinical Study. <i>J Baghdad College Dentistry</i> 2015;27:174-178.	Included

Appendix 4. Descriptive statistics of the re-analyzed provided data of Karamouzos et al.

Outcome	Teeth	Mean (SE)	95% CI	Range
Overall ΔE	416	2.80 (0.09)	2.62, 2.98	0.58-6.08
Outcome	Teeth (%)			
Teeth with $\Delta E > 3.7$	54 (13.0%)			
Teeth with $\Delta E > 3.7$/ patient	Patients (%)			
0	5 (20.0%)			
1	6 (24.0%)			
2	5 (20.0%)			
3	5 (20.0%)			
4	3 (12%)			
6	1 (4.0%)			

CI, confidence interval; SE, standard error.

Appendix 5. Re-analysis of the provided data of Karamouzos et al: generalized linear regression analysis on the treatment-induced color change (ΔE).

		Univariable			Multivariable	
Category	Group	β (95% CI)	P		β (95% CI)	P
Adhesive	Light-cured	Referent			Referent	
	Chemically-cured	0.41 (0.30,0.51)	<0.001		0.41 (0.30,0.51)	<0.001
Tooth	Central incisor	0.02 (-0.15,0.19)	0.82		0.02 (-0.15,0.19)	0.82
	Lateral incisor	0.31 (0.12,0.51)	0.001		0.31 (0.12,0.51)	0.001
	Canine	Referent			Referent	
	1 st premolar	0.30 (0.06,0.53)	0.01		0.30 (0.06,0.53)	0.01
Jaw	Maxilla	-0.11 (-0.31,0.10)	0.31		NT	
	Mandible	Referent				
Side	Right	-0.02 (-0.15,0.11)	0.76		NT	
	Left	Referent				

β , unstandardized regression coefficient; CI, confidence interval; NT, not tested.

Appendix 6. Re-analysis of the provided data of Karamouzos et al: generalized binary regression analysis on the number of teeth with treatment-induced color changes that are clinically visible ($\Delta E \geq 3.7$).

Category	Group	Univariable		Multivariable	
		RR (95% CI)	P	RR (95% CI)	P
Adhesive	Light-cured	Referent		Referent	
	Chemically-cured	2.60 (1.58,4.27)	<0.001	2.58 (1.58,4.23)	<0.001
Tooth	Central incisor	1.86 (0.95,3.62)	0.07	1.87 (0.95,3.66)	0.07
	Lateral incisor	3.00 (1.42,6.35)	0.004	2.97 (1.41,6.23)	0.004
	Canine	Referent		Referent	
	1 st premolar	1.86 (0.76,4.55)	0.18	1.87 (0.76,4.58)	0.17
Jaw	Maxilla	0.93 (0.46,1.89)	0.84	NT	
	Mandible	Referent			
Side	Right	1.35 (0.85,2.15)	0.21	NT	
	Left	Referent			

CI, confidence interval; RR, relative risk; NT, not tested.

Appendix 7. Further details on the present review.

Author contributions

RK, SNP, and TE conceived the idea. RK wrote the first draft of the protocol. RK, SNP, and TE revised the protocol. SNP performed the literature searches and extracted search hits. RK did screening by title, study selection by abstract and full-text, data extraction, and assessment of the risk of bias in duplicate, while SNP checked all procedures independently by replication and TE resolved any conflicts that arose. SNP handled communications with trialists and performed the statistical analysis. RK wrote the first draft of the manuscript. RK, SNP, and TE assisted in the interpretation of the results and revised the manuscript draft. SNP submitted the manuscript and is the guarantor and responsible for the accuracy of the data and for future updates of the review.

Data sharing

The full dataset from this study will be uploaded and will be freely available through Zenodo upon acceptance. The individual patient data provided by Dr. Karamouzos have not been uploaded and can be requested by Dr. Karamouzos himself.

Post hoc changes to the protocol

- Outcome selection: we had initially planned to include the overall color changes ΔE as primary outcome, the three segregated parameters (ΔL^* , Δa^* , and Δb^*) as secondary outcomes, and subjective assessments of color as secondary outcome. Due to the small number of contributing studies and the fact that the ΔL^* , Δa^* , and Δb^* are integrated in the ΔE , which is more clinically relevant, the three parameters ΔL^* , Δa^* , and Δb^* were not included finally and we focused on the primary outcome. The secondary outcome of subjective color assessment was not reported from any included studies.
- Additional analysis: a sensitivity analysis was conducted post hoc to alleviate the extreme heterogeneity seen in a meta-analysis. This was planned to be conducted in cases of high heterogeneity.
- Analyses of reporting biases were planned in protocol, but could not be conducted as < 10 studies were finally included.
 - Several subgroup analyses were planned in protocol, but could not be conducted as < 5 studies were finally included.